

Drug name	Recently used <input checked="" type="checkbox"/>	Drug name	Recently used <input checked="" type="checkbox"/>
MDMA/Ecstasy pills (non-prescribed)		PMA	
MDMA/Ecstasy powder (non-prescribed)		PMMA	
MDMA/Ecstasy crystal (non-prescribed)		Amyl nitrite/alkyl nitrite (poppers, jungle juice)	
MDMA/Ecstasy capsules (non-prescribed)		Nitrous oxide (nangs, bulbs)	
Methamphetamine powder (speed)		Cannabis and/or cannabinoid-related products	Prescribed/non-prescribed
Methamphetamine base (paste)		Tobacco (this includes tobacco that has been mixed with cannabis)	
Crystal methamphetamine (ice/shabu)		E-cigarettes	Prescribed/non-prescribed
Pharmaceutical stimulants (e.g., Ritalin/dexamfetamine/modafinil, lisdexamfetamine)	Prescribed/non-prescribed	Nicotine pouches	
Babazine		Alcohol	
Cocaine		Heroin	
LSD (acid)		Codeine	Prescribed/non-prescribed
Hallucinogenic mushrooms/psilocybin (non-prescribed)		Pharmaceutical opioids	Prescribed/non-prescribed

Kava		Benzodiazepines (e.g., Valium, Diazepam, Xanax, Kalma)	Prescribed/non-prescribed
MDA		Steroids (non-prescribed)	
Ketamine (special K) (non-prescribed)		Antipsychotics (e.g., Seroquel) (non-prescribed)	
GHB/GBL/1,4-BD (liquid E)		Unknown substance	
DMT			

## Past Six-Month Drug Use Service Utilisation

<b>SERVICE/PROFESSIONAL</b>	
1	<b>GP</b>
2	<b>Emergency Department</b>
3	<b>Hospital admission (inpatient)</b>
4	<b>Medical tent (at festival/rave)</b>
	<b>Dentist</b>
5	<b>Drug and alcohol counsellor</b>
6	<b>Hospital as outpatient</b>
7	<b>Specialist doctor (not including psychiatrists)</b>
9	<b>Other health professional (i.e. chiropractor, nurse, physiotherapist)</b>
10	<b>Ambulance attendance</b>
19	<b>Pharmacy</b>
13	<b>Psychologist</b>
11	<b>Psychiatrist</b>
15	<b>Needle and syringe program</b>
16	<b>Peer-based harm reduction service (DanceWize, Save-A-Mate)</b>
17	<b>Other harm reduction service</b>
18	<b>Other (specify)</b>

## New Psychoactive Substances (NPS)

No.	Drug name
0	2C - any (e.g., 2C-I, 2C-B)
1	2-fluoro-2-oxo PCE (CanKET, 2F-NENDCK)
2	2-fluorodeschloroketamine (2-FDCK)
3	3 CI-PCP/4CI-PCP
4	3-chloromethcathinone (3-CMC)
5	3F-2-oxo PCE (Fluorexetamine)
6	3-HO-PCP/4-HO-PCP
7	3-MeO-PCP/4- MeO-PCP
8	3-methylmethcathinone (3-MMC)
9	4-chloromethcathinone (4-CMC)
10	4-FA
11	4F-phenibut
12	5-MeO-DMT
13	8-aminoclonazepam
14	Alpha-PHP
15	Alpha-PVP
16	Ayahuasca
17	Bromazolam
18	Clonazolam
19	dimethylpentylone (N,N-dimethylpentylone, DMP)
20	DOX (e.g. DOB, DOC, DOI, DOM)
21	Etizolam
22	Flualprazolam
23	Flubromazepam
24	Glaucine
25	Kratom/ mitragynine
26	MDPV
27	Mephedrone
28	Methcathinone

- 29 Methylone
- 30 Methoxetamine (MXE)
- 31 Mescaline
- 32 NBOH (e.g. 25I, 25B)
- 33 NBOMe (e.g., 25I, 25B, 25C other)
- 34 N-ethylpentylone (ephylone)
- 35 N-ethyl hexedrone
- 36 N-ethylbutylone (eutylone)
- 37 Pentylone
- 38 Phenibut
- 39 Phenazolam
- 40 Salvia divinorum
- 41 Synthetic cannabinoids (e.g. ADB-BUTINACA, 4F-MDMB-BUTICA, FUB-AM)
- 42 Tiletamine
- 43 Tuci (also known as tusi, tucibi, pink cocaine)
- 44 Xylazine
- 45 Other 'new' drugs that mimic the effect of opioids (e.g. acetylfentanyl, nitazenes)
- 46 Other 'new' drugs that mimic the effect of ecstasy
- 47 Other 'new' drugs that mimic the effects of amphetamines or cocaine
- 48 Other 'new' drugs that mimic the effects of psychedelic drugs like LSD
- 49 Other 'new' drugs that mimic the effects of benzodiazepines
- 50 Other 'new' drugs that mimic the effects of dissociatives like ketamine
- Other new and emerging psychoactive substances

## Scale Sheet

### **K10 – LAST 4 WEEKS**

*These questions are related to how you have been feeling over the last 4 weeks. Remember, any information you provide is completely confidential.*

1. None of the time
2. A little of the time
3. Some of the time
4. Most of the time
5. All of the time

### **Ecstasy/Meth SDS – LAST 6 MONTHS**

*This section is about how you felt about your ecstasy and/or methamphetamine use in the last 6 months.*

1. Never or almost never
2. Sometimes
3. Often
4. Always or nearly always

### **AUDIT – LAST 12 MONTHS**

*These questions are related to your use of alcohol in the last year. Remember, any information you provide is completely confidential.*

1. Never
2. Less than monthly
3. Monthly
4. Weekly
5. Daily or almost daily

### **STIGMA**

*In the past six months, to what extent do you think healthcare workers have treated you poorly because of your use of illicit drugs?*

0. Never
1. Rarely
2. Sometimes
3. Often
4. Always
5. Haven't seen a specialist alcohol and other drug healthcare worker in the past 6 months

## Modes of Purchasing

### **Social networking and messaging**


**apps:** Refers to a type of online community such as Facebook or Instagram; messaging apps such as Whatsapp and snapchat; and dating apps like grindr and tinder.



**Surface web:** Any website accessible without the need for any special connection routing with a '.com' address.



**Dark web:** Any website only accessible via The Onion Router (TOR) with a '.onion' URL (e.g. Dark Market, White House Market, Monopoly Market, Cannazon).



**Face-to-Face:** Refers to purchasing illicit drugs in person.



## Risk Communication

### Example drug warnings:

#### **NSW Health has issued a drug warning:**

Fentanyl and acetyl fentanyl are circulating in Sydney and may be sold as methamphetamine.

These drugs are dangerous and can cause overdose and death.

Look out for unexpected effects such as drowsiness, slow breathing, loss of consciousness

Stay safe:

- Ring 000 if you or anyone else experiences these symptoms
- Don't use alone
- Carry naloxone
- Test your drugs the best way you can
- 

#### **Drug Warning**



Light blue “Lego” MDMA pills circulating in NSW

- Up to 3x common strength
- Dose may be variable in pill

Remember:

- Dropping in the heat is riskier
- High temps increase the neurotoxic effects of MDMA
- If you're planning on taking don't peak in the heat

Look out for:

- Excessive sweating or stopping sweating when active/hot
- Uncontrolled repetitive movements
- Confusion or agitation

Get your mates help if they are in trouble

Stay safe

- Stay as hydrated and as cool as possible
- Avoid mixing and don't use alone
- Seek medical help ASAP or call 000 if unwell



## Nitrous Oxide

### Nitrous oxide product size examples

Bulbs ~8g



Bottles  $\leq 1000\text{g}$   
(common sizes = 0.95L/~580g; 1L/~615g)



Bottles >1000g  
(common sizes = 2.2L/~1100g; 3.3L/~2100g)



## Cocaine Forms

Picture 1



Picture 2



Picture 3



Picture 4

