

**Mortality among people with substance use disorders: A toolkit for classifying major causes of death**

**Thomas Santo Jr., Chrianna Bharat, Samantha Colledge-Frisby, Agata Chrzanowska, Nicola Man, Lauren Moran, Emma Torrens, & Louisa Degenhardt**

**NDARC Technical Report No. 301**

**Recommended citation:** Santo, Thomas, Bharat, C., Colledge-Frisby, S., Chrzanowska, A., Man, N., Moran, L., Torrens, E., & Degenhardt, L. (2022). *Mortality among people with substance use disorders: A toolkit for classifying major causes of death*. Sydney: National Drug and Alcohol Research Centre, UNSW Sydney.

**ISBN: 978-0-7334-4032-8**

**©NATIONAL DRUG AND ALCOHOL RESEARCH CENTRE, UNSW SYDNEY, 2022**

**This work is copyright. You may download, display, print and reproduce this material in unaltered form only (retaining this notice) for your personal, non-commercial use or use within your organisation.**

**All other rights are reserved. Requests and enquiries concerning reproduction and rights should be addressed to the information manager, National Drug and Alcohol Research Centre, UNSW Sydney, NSW 2052, Australia.**

## Overview

- Some people with substance use disorders, particularly people who inject drugs and people with cocaine<sup>1</sup>, amphetamine<sup>2</sup>, or opioid dependence<sup>3</sup>, are at elevated risk of mortality. These elevated risks tend to be concentrated across a few key domains. Some of those are related to the manner in which drugs are taken (e.g. contraction of blood borne viral infections such as HIV through sharing of injection equipment); others may be related to the effects of drug and alcohol use (e.g. liver disease) and others still as relatively direct consequences of acute drug intoxication (e.g. accidental drug overdose).
- Classifying deaths into appropriate categories is not a straightforward task. Multiple iterations of the World Health Organization's (WHO) International Classification of Diseases exist, each of which defines deaths slightly differently. Some codes are not readily interpretable as belonging to a distinct grouping. Hundreds of codes exist, so the volume of categories may present challenges.
- This bulletin provides an overview of some key references that provide guidance on classifying major causes of death that occur among people who use illicit drugs. It is intended to make categorization of deaths among illicit drug users an easier process, and perhaps one that is undertaken more consistently across studies.
- It provides groupings for both the International Classification of Diseases Version 9 (ICD-9) and Version 10 (ICD-10). The comparable codes in each are listed alongside each other. There may be differences in the deaths captured due to the version change, and this should be noted when analysing data across time.
- The source of each of these groupings is provided alongside each for ease of reference.
- Feedback on these groupings or suggestions for further groupings are welcomed.
- This is an update of a technical report from 2009<sup>4</sup>

**Table 1: Definitions for major causes of death**

Cause of death category	ICD-9	ICD-10	Sources
<b>All injury and poisoning (excl. alcohol-related)</b>	304, 305.2-305.9, E800-E869, E880-E929, E950-E999	F11-F16, F19, F55, V00-X99, Y00-Y39, Y85-Y87, Y89	CHO injury and poisoning indicator ( <a href="#">Link</a> ) <ul style="list-style-type: none"> <li>- NB: F codes added to ICD-10 to include drug related deaths in circumstances of dependence (likewise 300 codes for ICD-9) and to encapsulate all deaths in subcategories below</li> <li>- Does not include accidental or intentional poisoning by exposure to alcohol (X45, X65, Y15)</li> </ul>
<b>Drug-induced deaths</b> <i>Two definitions are provided. The first is a narrower definition and captures codes specifically related to drug use disorders and poisonings.</i>	304, 305.2-305.9, E850-E858, E950.0-E950.5, E962.0, E980.0-E980.5	F11- F16, F19, F55, X40- X44, X60-X64, X85, Y10-Y14	Santo, T., Clark, B., Hickman, M., Grebely, J., Campbell, G., Sordo, L., ... & Degenhardt, L. (2021). Association of opioid agonist treatment with all-cause mortality and specific causes of death among people with opioid dependence: a systematic review and meta-analysis. JAMA psychiatry <sup>5</sup>
<i>The second definition<sup>1</sup>, is broader and captures drug-related diseases in addition to drug use disorders and poisonings. Note, there are several ICD-10 codes in this definition for which there is no corresponding ICD-9 code due to lack of mention of drug-induced. These include D, E, F18, G, I, J, L, and M. <sup>2</sup></i>	292, 304, 305.2-305.9, 790.99, E850-E858, E950.0-E950.5, E962.0, E980.0-E980.5	<i>Underlying cause of death:</i> F11.0-F11.5, F11.7-F11.9, F12.0-F12.5, F12.7-F12.9, F13.0-F13.5, F13.7-F13.9, F14.0-F14.5, F14.7-F14.9, F15.0-F15.5, F15.7-F15.9, F16.0-F16.5, F16.7-F16.9, F18.0-F18.5, F18.7-F18.9, F19.0-F19.5, F19.7-F19.9, D52.1, D59.0, D59.2, D61.1, D64.2, E06.4, E16.0, E23.1, E24.2, E27.3, E66.1, G21.1, G24.0, G25.1, G25.4, G25.6, G44.4, G62.0, G72.0, I95.2, J70.2-J70.4, L10.5, L27.0, L27.1, M10.2, M32.0, M80.4, M81.4, M83.5, M87.1, R78.1-R78.5, X40-X44, X60-X64, X85 Y10-Y14	Chrzanowska, A., Man, N., Sutherland, R., Degenhardt, L. & Peacock, A. (2021). Methods for “Trends in drug-induced deaths in Australia, 1997-2019”. Drug Trends Bulletin Series. Sydney: National Drug and Alcohol Research Centre, UNSW Sydney. <sup>6</sup>  3303.0 - Causes of Death, Australia, 2016 ( <a href="#">Link</a> )
<b>Suicide</b>	E950-E959	X60- X84, Y87.0	CHO suicide indicator ( <a href="#">Link</a> )
Non-poisoning suicides	E950.6-E959	X66-X85, Y87.0	

Cause of death category	ICD-9	ICD-10	Sources
<b>Violence</b>	E960-E969	X85-Y09, Y87.1	CHO violence indicator ( <a href="#">Link</a> )
<b>Motor vehicle and transport accidents</b>	E810-E825, E929.0	V01-V99	CHO motor vehicle crash injury indicator ( <a href="#">Link</a> ) and CHO unintentional injuries indicators ( <a href="#">Link</a> ) - NB: modified to only the ICD-10 chapter 'Transport Accidents' including all codes in this chapter
<b>Falls/fires/burns/drownings</b>	E880-E886, E888; E890-E899; E910	W00-W19, W65-W74, X00-X09	CHO unintentional injuries indicators ( <a href="#">Link</a> )
<b>All liver-related</b>	070, 155, 570, 571, 572-573	B15–B19, B94.2, C22, I85.0, K70–K77, O98.4, P35.3	Amin J, Law M, Bartlett M, Kaldor J, Dore G. Causes of death after diagnosis of hepatitis B or hepatitis C infection: a large community-based linkage study. Lancet 2006; 368: 938–45. - Added obstetric and neonatal hepatitis codes from CHO viral hepatitis indicator: NB: I85.0 (Oesophageal varices with bleeding) added
Viral hepatitis	070	B15-B19, B94.2, I85.0, O98.4, P35.3	CHO potentially avoidable mortality table - ( <a href="#">Link</a> ) - NB: I85.0 (Oesophageal varices with bleeding) added
<b>All alcohol-related</b>	291, 303.0, 303.9, 305.0, 357.5, 425.5, 535.3, 571.0-571.3, E860.0-E860.2 <sup>2</sup>	E24.4, F10, G31.2, G62.1, G72.1, I42.6, K29.2, K85.2, K86.2, K70, K86.0, R78.0, X45, X65, Y15	- Causes of Death, Australia methodology, 2020 ( <a href="#">Link</a> ); - ICD9 translation assisted by Chikritzhs et al 2002 'Alcohol-related codes: Mapping ICD-9 to ICD-10' Technical Report from NDRI and Dept of Health, Western Australia ( <a href="#">Link</a> )
<b>Cancer</b>	140-208, 238.4, 238.7, 273.1	C00-C97, D45-D46, 47.1, D47.3-D47.5	CHO indicator for cancer ( <a href="#">Link</a> )
<b>Cardiovascular disease</b>	390-459	I00-I99, G45, G46	CHO cardiovascular disease indicator ( <a href="#">Link</a> )
<b>Chronic lower respiratory disease</b>	490, 491, 492, 493, 496	J40-J46	CHO chronic respiratory disease indicator ( <a href="#">Link</a> )
<b>Digestive disorders</b>	531-534, 540-543, 550-553, 574-577	K25-K28, K35-K38, K40-K46, K80-K83, K85-K86, K91.5	CHO potentially avoidable mortality table ( <a href="#">Link</a> )
<b>HIV-related</b>	042-044, 279.10	B20-B24	ABS definition in 3303.0 - Causes of Death, Australia, 2006, explanatory notes point 42 ( <a href="#">Link</a> )
<b>Influenza and pneumonia</b>	480-488	J09-J18	CHO indicator ( <a href="#">Link</a> )
<b>Injecting-related diseases</b>	038, 040.0, 040.8, 041.0, 041.1, 041.8, 041.9, 112.5, 112.81, 324.X, 394.0, 394.2, 394.9, 395, 396.8, 397.0, 421.X, 424.X, 451, 453.2, 453.3, 453.8, 453.9, 459.1,	A48.0, G06.0, G06.1, G06.2, L02.X, L03.X, L08.8, L08.9, L97, L98.4, L98.8, L98.9, M72.6, R02, B37.6, I33.0, I33.9, I34.0, I34.2, I34.8, I34.9, I35.X, I36.X, I37.X, I38, I39.X, T82.6, A40.X, A41.X, A49.1, A49.8, A49.9, B37.7, R57.2, R65.1, R65.9,	Colledge-Frisby, S., Jones, N., Larney, S., Peacock, A., Lewer, D., Brothers, T. D., Hickman, M., Farrell, M., & Degenhardt, L. (in press). The impact of opioid agonist treatment on hospitalisations or injecting-related diseases among an opioid dependent population: A retrospective data linkage study. Drug Alcohol Depend. <sup>7</sup>

Cause of death category	ICD-9	ICD-10	Sources
	459.81, 459.89, 459.9, 680.X, 681.X, 682.X, 686.8, 707.1, 707.8, 707.9, 709.8, 709.9, 711.0, 711.4, 711.9, 722.9, 728.86, 730.0, 730.1, 730.2, 730.8, 730.9, 746.00, 785.4, 785.52, 995.9, 995.90, 995.91, 996.61	M00.X, M86.X, M89.9, M46.2, M46.3, M46.4, I80, I82.2, I82.3, I82.8, I82.9, I87.0, I87.2, I87.8, I87.9, A48.8, A49.0, I26.9	
<b>Skin and soft tissue infections</b>	040.0, 324.X, 680.X, 681.X, 682.X, 686.8, 686.9, 707.1, 707.8, 707.9, 709.8, 709.9, 728.86, 785.4	A48.0, G06.0, G06.1, G06.2, L02.X, L03.X, L08.8, L08.9, L97, L98.4, L98.8, L98.9, M72.6, R02	Colledge-Frisby, S., Jones, N., Larney, S., Peacock, A., Lewer, D., Brothers, T. D., Hickman, M., Farrell, M., & Degenhardt, L. (in press). The impact of opioid agonist treatment on hospitalisations or injecting-related diseases among an opioid dependent population: A retrospective data linkage study. Drug Alcohol Depend. <sup>7</sup>
<b>Endocarditis</b>	112.81, 394.0, 394.2, 394.9, 395, 396.8, 397.0, 421.X, 424.X, 746.00, 996.61	B37.6, I33.0, I33.9, I34.0, I34.2, I34.8, I34.9, I35.X, I36.X, I37.X, I38, I39.X, T82.6	Colledge-Frisby, S., Jones, N., Larney, S., Peacock, A., Lewer, D., Brothers, T. D., Hickman, M., Farrell, M., & Degenhardt, L. (in press). The impact of opioid agonist treatment on hospitalisations or injecting-related diseases among an opioid dependent population: A retrospective data linkage study. Drug Alcohol Depend. <sup>7</sup>
<b>Sepsis and bacteraemia</b>	040.8, 041.1, 038, 038.0, 038.1, 038.10, 038.19, 038.2, 038.3, 038.4, 038.41, 038.8, 038.9, 041.0, 041.8, 041.9, 112.5, 785.52, 995.9, 995.90, 995.91	A40.X, A41.X, A49.1, A49.8, A49.9, B37.7, R57.2, R65.1, R65.9, A48.8, A49.0, I26.9	Colledge-Frisby, S., Jones, N., Larney, S., Peacock, A., Lewer, D., Brothers, T. D., Hickman, M., Farrell, M., & Degenhardt, L. (in press). The impact of opioid agonist treatment on hospitalisations or injecting-related diseases among an opioid dependent population: A retrospective data linkage study. Drug Alcohol Depend. <sup>7</sup>
<b>Septic arthritis</b>	711.0, 711.4, 711.9	M00.X	Colledge-Frisby, S., Jones, N., Larney, S., Peacock, A., Lewer, D., Brothers, T. D., Hickman, M., Farrell, M., & Degenhardt, L. (in press). The impact of opioid agonist treatment on hospitalisations or injecting-related diseases among an opioid dependent population: A retrospective data linkage study. Drug Alcohol Depend. <sup>7</sup>
<b>Osteomyelitis</b>	722.9, 730.0, 730.1, 730.2, 730.8, 730.9	M86.X, M89.9, M46.2, M46.3, M46.4	Colledge-Frisby, S., Jones, N., Larney, S., Peacock, A., Lewer, D., Brothers, T. D., Hickman, M., Farrell, M., & Degenhardt, L. (in press). The impact of opioid agonist treatment on hospitalisations or injecting-related diseases among an opioid dependent population: A retrospective data linkage study. Drug Alcohol Depend. <sup>7</sup>

Cause of death category	ICD-9	ICD-10	Sources
<b>Venous diseases</b>	451, 453.2, 453.3, 453.8, 453.9, 459.1, 459.81, 459.89, 459.9	I80, I82.2, I82.3, I82.8, I82.9, I87.0, I87.2, I87.8, I87.9	Colledge-Frisby, S., Jones, N., Larney, S., Peacock, A., Lewer, D., Brothers, T. D., Hickman, M., Farrell, M., & Degenhardt, L. (in press). The impact of opioid agonist treatment on hospitalisations or injecting-related diseases among an opioid dependent population: A retrospective data linkage study. <i>Drug Alcohol Depend.</i> <sup>7</sup>

**Table Notes:** <sup>1</sup> Provided by the Australian Bureau of Statistics; <sup>2</sup> Table 2 provides drug-induced death definitions by drug class and Table 3 provides drug-induced deaths definitions by intent; <sup>3</sup> No equivalent ICD-9 code for acute alcoholic pancreatitis

**Table 2. Drug-induced deaths by drug class**

<b>Drug class</b>	<b>ICD-9</b>	<b>ICD-10</b>
<b>Antiepileptic, sedative-hypnotic, and antiparkinsonism drugs</b>	304.1*, 305.4*	<i>UCOD</i> : Drug-induced death <i>MCOD</i> : T42.0-T42.8
<b>Opioids</b>	304.0*, 304.7*, 305.5*, E935.0-E935.2	<i>UCOD</i> : Drug-induced death <i>MCOD</i> : T40.0-T40.4, T40.6 <b>OR</b> <i>UCOD</i> : F11 <i>MCOD</i> : (any)
<b>Amphetamines &amp; other stimulants</b>	304.4*, 305.7*	<i>UCOD</i> : Drug-induced death <i>MCOD</i> : T43.6 <b>OR</b> <i>UCOD</i> : F15 <i>MCOD</i> : (any)
<b>Antidepressants</b>	305.8, E939.0	<i>UCOD</i> : Drug-induced death <i>MCOD</i> : T43.0-T43.2
<b>Antipsychotic and neuroleptics</b>	E939.3	<i>UCOD</i> : Drug-induced death <i>MCOD</i> : T43.3-T43.5
<b>Non-opioid analgesic, antipyretics, and antirheumatics</b>	965.1-965.9, E935.3-E935.9	<i>UCOD</i> : Drug-induced death <i>MCOD</i> : T39.0-T39.9
<b>Alcohol</b>	291, 303, 305.0*	<i>UCOD</i> : Drug-induced death <i>MCOD</i> : T51.0, T51.9
<b>Cannabis derivatives</b>	304.3*, 305.2*	<i>UCOD</i> : Drug-induced death <i>MCOD</i> : T40.7 <b>OR</b> <i>UCOD</i> : F12 <i>MCOD</i> : (any)
<b>Cocaine</b>	304.2*, 305.6*	<i>UCOD</i> : Drug-induced death <i>MCOD</i> : T40.5 <b>OR</b> <i>UCOD</i> : F14 <i>MCOD</i> : (any)

**Table Notes:** *UCOD*, underlying/primary cause of death; *MCOD*, multiple causes of death;

**Table 3. Drug-induced deaths by intent**

Cause of death category	ICD-9	ICD-10
<b>Accidental drug-induced deaths</b>	E850-E858	<p><i>UCOD</i>: X40-X44</p> <p><b>OR</b></p> <p><i>UCOD</i> (Prior to 2013<sup>a</sup>):            F11.0-F11.5, F11.7-F11.9,            F12.0-F12.5, F12.7-F12.9,            F13.0-F13.5, F13.7-F13.9,            F14.0-F14.5, F14.7-F14.9,            F15.0-F15.5, F15.7-F15.9,            F16.0-F16.5, F16.7-F16.9,            F18.0-F18.5, F18.7-F18.9,            F19.0-F19.5, F19.7-F19.9,  <i>MCOD</i>: X40-X44</p>
<b>Accidental opioid-induced deaths</b>	E850.0	<p><i>UCOD</i>: X42</p> <p><i>MCOD</i>: T40.0-T40.4 or T40.6            [Excl. if T40.6 with (T40.5, T40.8 or T40.9)]</p> <p><b>OR</b></p> <p><i>UCOD</i>: X44</p> <p><i>MCOD</i>: T40.0-T40.4 or T40.6</p> <p><b>OR</b></p> <p><i>UCOD</i>: X40, X41, X43</p> <p><i>MCOD</i>: (T40.0-T40.4 or T40.6) AND no other T-codes present</p> <p><b>OR</b></p> <p><i>UCOD</i> (Prior to 2013<sup>a</sup>):            F11.0-F11.5, F11.7-F11.9,            F12.0-F12.5, F12.7-F12.9,            F13.0-F13.5, F13.7-F13.9,            F14.0-F14.5, F14.7-F14.9,            F15.0-F15.5, F15.7-F15.9,            F16.0-F16.5, F16.7-F16.9,            F18.0-F18.5, F18.7-F18.9,            F19.0-F19.5, F19.7-F19.9,</p> <p><b>AND</b></p> <p>{ <i>MCOD</i>: X42 with (T40.0-T40.4 or T40.6) [Excl. if T40.6 with (T40.5, T40.8 or T40.9)]</p> <p><b>or</b></p> <p><i>MCOD</i>: X44 with (T40.0-T40.4 or T40.6)</p> <p><b>or</b></p> <p><i>MCOD</i>: X40, X41, X43 with (T40.0-T40.4 or T40.6) AND no other T-codes present }</p>
<b>Intentional drug-induced deaths</b>	E950.0-E950.5	<p><i>UCOD</i>: X60-X64</p> <p><b>OR</b></p> <p><i>UCOD</i> (Prior to 2013<sup>a</sup>):            F11.0-F11.5, F11.7-F11.9,            F12.0-F12.5, F12.7-F12.9,            F13.0-F13.5, F13.7-F13.9,            F14.0-F14.5, F14.7-F14.9,            F15.0-F15.5, F15.7-F15.9,            F16.0-F16.5, F16.7-F16.9,            F18.0-F18.5, F18.7-F18.9,            F19.0-F19.5, F19.7-F19.9,  <i>MCOD</i>: X60-X64}</p>



<p><b>Intentional opioid-induced deaths</b></p>	<p>No corresponding code<sup>1</sup></p>	<p><i>UCOD</i>: X62  <i>MCOD</i>: T40.0-T40.4 or T40.6  [Excl. if T40.6 with (T40.5, T40.8 or T40.9)]  <b>OR</b>  <i>UCOD</i>: X64  <i>MCOD</i>: T40.0-T40.4 or T40.6  <b>OR</b>  <i>UCOD</i>: X60, X61, X63  <i>MCOD</i>: (T40.0-T40.4 or T40.6) AND no other T-codes present  <b>OR</b>  (Prior to 2013)  <i>UCOD</i>:  F11.0-F11.5, F11.7-F11.9,  F12.0-F12.5, F12.7-F12.9,  F13.0-F13.5, F13.7-F13.9,  F14.0-F14.5, F14.7-F14.9,  F15.0-F15.5, F15.7-F15.9,  F16.0-F16.5, F16.7-F16.9,  F18.0-F18.5, F18.7-F18.9,  F19.0-F19.5, F19.7-F19.9,  <b>AND</b>  {<i>MCOD</i>: X42 with (T40.0-T40.4 or T40.6) [Excl. if T40.6 with (T40.5, T40.8 or T40.9)]  <b>or</b>  <i>MCOD</i>: X44 with (T40.0-T40.4 or T40.6)  <b>or</b>  <i>MCOD</i>: X40, X41, X43 with (T40.0-T40.4 or T40.6) AND no other T-codes present}</p>
<p><b>Undetermined drug-induced deaths</b></p>	<p>E980.0-E980.5</p>	<p><i>UCOD</i>: Y10-Y14  <b>OR</b>  (Prior to 2013^)  <i>UCOD</i>:  F11.0-F11.5, F11.7-F11.9,  F12.0-F12.5, F12.7-F12.9,  F13.0-F13.5, F13.7-F13.9,  F14.0-F14.5, F14.7-F14.9,  F15.0-F15.5, F15.7-F15.9,  F16.0-F16.5, F16.7-F16.9,  F18.0-F18.5, F18.7-F18.9,  F19.0-F19.5, F19.7-F19.9,  <i>MCOD</i>: Y10-Y14</p>
<p><b>Undetermined opioid-induced deaths</b></p>	<p>No corresponding code<sup>1</sup></p>	<p><i>UCOD</i>: Y12  <i>MCOD</i>: T40.0-T40.4 or T40.6  [Excl. if T40.6 with (T40.5, T40.8 or T40.9)]  <b>OR</b>  <i>UCOD</i>: Y14  <i>MCOD</i>: T40.0-T40.4 or T40.6  <b>OR</b>  <i>UCOD</i>: Y10, Y11, Y13  <i>MCOD</i>: (T40.0-T40.4 or T40.6) AND no other T-codes present  <b>OR</b></p>

	<p><i>UCOD: (Prior to 2013<sup>^</sup>):</i>  F11.0-F11.5, F11.7-F11.9,  F12.0-F12.5, F12.7-F12.9,  F13.0-F13.5, F13.7-F13.9,  F14.0-F14.5, F14.7-F14.9,  F15.0-F15.5, F15.7-F15.9,  F16.0-F16.5, F16.7-F16.9,  F18.0-F18.5, F18.7-F18.9,  F19.0-F19.5, F19.7-F19.9,  <b>AND</b>  { MCODE: Y12 with (T40.0-T40.4 or T40.6) [Excl. if T40.6 with (T40.5,  T40.8 or T40.9)]  <b>or</b>  MCODE: Y14 with (T40.0-T40.4 or T40.6)  <b>or</b>  MCODE: Y10, Y11, Y13 with (T40.0-T40.4 or T40.6) AND no other T-  codes present }</p>
--	--

**Table Notes:** UCOD, underlying/primary cause of death; MCODE, multiple causes of death;

<sup>1</sup>No ICD-9 codes directly match ICD-10 undetermined and intentional opioid deaths. However, since ICD-9 has no contributing cause categories, all ICD-9 opioid-related deaths are technically undetermined, including 304.0\*, 304.7\*, and 305.5\*. Researchers may consider defining all ICD-9 non-accidental opioid-related deaths as undetermined (incl. intentional) but should be reported separately from deaths coded **using** ICD-10 in this category.

<sup>^</sup>Prior to the 2013 reference year, the 2006 version of the ICD-10 was the most recent version used for coding deaths, with the exception of two updates after 2006. The first update was in 2007 related to the use of mental and behavioural disorders due to psychoactive substance use, acute intoxication (F10.0, F11.0....F19.0) as an underlying cause of death. If the acute intoxication initiated the train of morbid events, it is now assigned an external accidental poisoning code (X40-X49) corresponding to the type of drug used. For example, if the death had been due to alcohol intoxication, the underlying cause before the update was F10.0, and after the update the underlying cause is X45, with poisoning code T51.9.

## References

1. Peacock A, Tran LT, Larney S, Stockings E, Santo Jr T, Jones H, et al. All-cause and cause-specific mortality among people with regular or problematic cocaine use: a systematic review and meta-analysis. *Addiction*. 2021;116:725–42.
2. Stockings E, Tran LT, Santo Jr T, Peacock A, Larney S, Santomauro D, et al. Mortality among people with regular or problematic use of amphetamines: a systematic review and meta-analysis. *Addiction*. 2019;114:1738–50.
3. Larney S, Tran LT, Leung J, Santo T, Santomauro D, Hickman M, et al. All-Cause and Cause-Specific Mortality Among People Using Extramedical Opioids. *JAMA Psychiatry* [Internet]. 2019; Available from: <https://dx.doi.org/10.1001/jamapsychiatry.2019.4170>
4. Randall D, Roxburgh A, Gibson A, Degenhardt L. Mortality among people who use illicit drugs: A toolkit for classifying major causes of death. *Syd Natl Drug Alcohol Res Cent Univ NSW*. 2009;
5. Santo T Jr, Clark B, Hickman M, Grebely J, Campbell G, Sordo L, et al. Association of Opioid Agonist Treatment With All-Cause Mortality and Specific Causes of Death Among People With Opioid Dependence: A Systematic Review and Meta-analysis. *JAMA Psychiatry* [Internet]. 2021 [cited 2021 Jun 8]; Available from: <https://doi.org/10.1001/jamapsychiatry.2021.0976>
6. Trends In Drug-Induced Deaths In Australia, 1997-2019 [Internet]. Sydney: National Drug and Alcohol Research Centre, UNSW Sydney; 2021. Available from: [http://handle.unsw.edu.au/1959.4/unsworks\\_75253](http://handle.unsw.edu.au/1959.4/unsworks_75253)
7. Colledge-Frisby S, Jones N, Larney S, Peacock A, Lewer D, Brothers TD, et al. The impact of opioid agonist treatment on hospitalisations or injecting-related diseases among an opioid dependent population: A retrospective data linkage study. *Drug Alcohol Depend*.