

AUSTRALIAN INSTITUTE OF HEALTH AND WELFARE  
NATIONAL PERINATAL STATISTICS UNIT  
AND  
THE FERTILITY SOCIETY OF AUSTRALIA  
ASSISTED CONCEPTION SERIES  
Number 4

# **Assisted conception Australia and New Zealand 1997**

**Tara Hurst  
Esther Shafir  
and  
Paul Lancaster**

AIHW National Perinatal Statistics Unit  
Sydney, 1999

AIHW cat. no. PER 10

© Australian Institute of Health and Welfare 1999

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced without prior written permission from the Australian Institute of Health and Welfare. Requests and enquiries concerning reproduction and rights should be directed to the Head, Communication and Public Affairs, Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601.

This is the fourth publication of the Australian Institute of Health and Welfare National Perinatal Statistics Unit's Assisted Conception Series. A complete list of the AIHW National Perinatal Statistics Unit's publications is available from the Publications Unit, Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601, or via the NPSU web site (<http://www.aihw.gov.au/npsu/>)

ISSN 1038 7234  
ISBN 1 74024 009 X

### **Suggested citation**

Hurst T, Shafir E & Lancaster P 1999. Assisted conception Australia and New Zealand 1997. AIHW cat. no. PER 10. Sydney: Australian Institute of Health and Welfare National Perinatal Statistics Unit (Assisted Conception Series no. 4).

### **Australian Institute of Health and Welfare**

Board Chair  
Professor Janice Reid

Director  
Dr Richard Madden

Published by Australian Institute of Health and Welfare National Perinatal Statistics Unit  
Printed by Ausdoc on Demand

# Contents

List of figures.....	v
List of tables.....	vi
Acknowledgments.....	x
Collaborating IVF and GIFT units.....	xi
Abbreviations.....	xiii
Highlights.....	1
1 Introduction.....	3
1.1 Variations in use of assisted conception.....	3
2 Treatment cycles and pregnancy rates.....	5
2.1 IVF and GIFT treatment cycles and pregnancy rates in 1997.....	6
3 IVF pregnancies.....	8
3.1 Maternal and paternal characteristics.....	8
3.1.1 Place of residence.....	8
3.1.2 Parental age.....	8
3.1.3 Previous pregnancies.....	9
3.1.4 Duration of infertility.....	9
3.1.5 Causes of infertility.....	9
3.2 Management of IVF pregnancies.....	10
3.2.1 Ovarian stimulation.....	10
3.2.2 Treatment cycle in which pregnancy occurred.....	11
3.2.3 Number of oocytes collected.....	11
3.2.4 Number of embryos transferred.....	11
3.2.5 Donor or frozen gametes and embryos.....	12
3.2.6 Microinsemination.....	13
3.2.7 Drugs used in luteal phase of pregnancy.....	13
3.3 Outcome of pregnancy.....	13
3.3.1 Maternal deaths.....	13
3.3.2 Maternal age and outcome of pregnancy.....	13
3.3.3 Spontaneous abortion.....	14
3.3.4 Ectopic pregnancy.....	14
3.3.5 Heterotopic pregnancies.....	15
3.3.6 Selective reduction of fetuses.....	15
3.3.7 Complications of pregnancy.....	15

3.3.8	Viable pregnancies of at least 20 weeks' gestation.....	16
3.3.9	Multiple pregnancies.....	17
3.3.10	Method of delivery .....	19
3.3.11	Sex of infants.....	19
3.3.12	Birthweight .....	20
3.3.13	Perinatal mortality .....	20
3.3.14	Congenital malformations .....	22
4	GIFT pregnancies.....	23
4.1	Maternal and paternal characteristics.....	23
4.1.1	Place of residence .....	23
4.1.2	Parental age.....	23
4.1.3	Previous pregnancies.....	24
4.1.4	Duration and causes of infertility .....	24
4.2	Management of GIFT pregnancies .....	24
4.2.1	Ovarian stimulation.....	24
4.2.2	Number of oocytes collected and transferred.....	25
4.2.3	Drugs used in luteal phase of pregnancy .....	25
4.3	Outcome of pregnancy .....	26
4.3.1	Maternal deaths.....	26
4.3.2	Spontaneous abortion and ectopic pregnancy.....	26
4.3.3	Heterotopic pregnancies .....	26
4.3.4	Complications of pregnancy.....	26
4.3.5	Viable pregnancies of at least 20 weeks' gestation.....	26
4.3.6	Multiple pregnancies.....	27
4.3.7	Method of delivery .....	27
4.3.8	Sex of infants.....	27
4.3.9	Infant's birthweight .....	27
4.3.10	Perinatal mortality .....	27
4.3.11	Congenital malformations .....	28
5	Tables.....	29
6	Bibliography .....	68
	Appendix 1: Definitions .....	74
	Appendix 2: Notification form .....	75

# List of figures

Figure 1: Ratio of treatment cycles to number of women in reproductive age group, 1997 .....	4
Figure 2: Assisted conception pregnancies by method of conception, 1990–1997.....	5
Figure 3: Outcome of IVF pregnancies by year of conception, 1979–1997 .....	8
Figure 4: Place of parental residence, IVF pregnancies, 1979–1997 .....	9
Figure 5: Duration of infertility, IVF pregnancies, 1979–1997 .....	10
Figure 6: Causes of infertility, IVF pregnancies, 1979–1997 .....	10
Figure 7: Drugs used to stimulate ovulation, IVF pregnancies, 1979–1997 .....	11
Figure 8: Mean number of oocytes collected by laparoscopy or ultrasound guidance, IVF and GIFT pregnancies, 1987–1997 .....	12
Figure 9: Percentage distribution of number of embryos transferred, IVF pregnancies, 1979–1997.....	12
Figure 10: Mean number of embryos/oocytes transferred, IVF and GIFT pregnancies, 1979–1997.....	13
Figure 11: Incidence of ectopic pregnancy and spontaneous abortion, IVF pregnancies, 1979–1997.....	14
Figure 12: Spontaneous abortions by maternal age groups, IVF and GIFT pregnancies, 1979–1997.....	15
Figure 13: Incidence of preterm birth in singleton IVF pregnancies, 1979–1997.....	17
Figure 14: Incidence of twin pregnancy, IVF and GIFT, 1979–1997.....	18
Figure 15: Incidence of triplet pregnancy, IVF and GIFT, 1979–1997 .....	18
Figure 16: Multiple pregnancies after assisted conception, Australia and New Zealand, 1996–1997.....	19
Figure 17: Caesarean birth rates, singleton and multiple IVF pregnancies, 1979–1997 .....	20
Figure 18: Birthweight of IVF and GIFT births, 1979–1997 .....	21
Figure 19: Incidence of low birthweight in singleton IVF births, 1979–1997.....	21
Figure 20: Perinatal mortality in singleton and multiple IVF births, 1979–1997.....	22
Figure 21: Outcome of GIFT pregnancies by year of conception, 1985–1997 .....	23
Figure 22: Causes of infertility, GIFT pregnancies, 1985–1997 .....	24
Figure 23: Percentage distribution of number of oocytes transferred, GIFT pregnancies, 1985–1997.....	25
Figure 24: Perinatal mortality in singleton and multiple GIFT births, 1985–1997 .....	28

# List of tables

Table 1:	Use of assisted conception to treat infertility, selected States, Australia and New Zealand, 1997 .....	29
Table 2:	Viable pregnancy rates for all techniques of assisted conception, 1997 .....	29
Table 3:	Oocyte retrieval cycles for IVF, ICSI and GIFT, by maternal age, cause of infertility, and drugs used to stimulate ovulation, 1997 .....	30
Table 4:	Embryo transfer cycles for IVF, ICSI and GIFT, by number of embryos or oocytes transferred, 1997.....	30
Table 5:	Number of transfer cycles after cryopreservation, by maternal age, cause of infertility, and number of embryos transferred, 1997.....	31
Table 6:	IVF pregnancies after transfer of fresh embryos to uterus, numbers and pregnancy rates in each IVF unit, 1997 .....	32
Table 7:	IVF pregnancies after ICSI, numbers and pregnancy rates in each IVF unit, 1997 .....	33
Table 8:	Pregnancy rates by type of transfer and technique of sperm collection, ICSI pregnancies, 1997 .....	34
Table 9:	IVF pregnancies after use of donor oocytes, numbers and pregnancy rates in each IVF unit, 1997 .....	34
Table 10:	GIFT pregnancies, numbers and pregnancy rates in each IVF unit, 1997 .....	35
Table 11:	IVF pregnancies after embryo freezing but without ICSI, numbers and pregnancy rates in each IVF unit, 1997 .....	36
Table 12:	IVF pregnancies after ICSI and embryo freezing, numbers and pregnancy rates in each IVF unit, 1997.....	37
Table 13:	Treatment related to embryo freezing in each IVF unit, 1997 .....	38
Table 14:	Numbers and outcomes of IVF pregnancies by year of conception, 1979–1997.....	38
Table 15:	Place of parental residence, IVF pregnancies, 1979–1997 .....	39
Table 16:	Maternal ages, IVF pregnancies, 1979–1997.....	39
Table 17:	Paternal ages, IVF pregnancies, 1979–1997 .....	39
Table 18:	Previous pregnancies for pregnant women, IVF pregnancies, 1979–1997 .....	40
Table 19:	Duration of infertility, IVF pregnancies, 1979–1997 .....	40
Table 20:	Outcome of pregnancy by duration of infertility, IVF pregnancies, 1996–1997.....	40
Table 21:	Causes of infertility, IVF pregnancies, 1979–1997 .....	41
Table 22:	Outcome of pregnancy by causes of infertility, IVF pregnancies, 1996–1997 .....	41
Table 23:	Drugs used to stimulate ovulation, IVF pregnancies, 1979–1997 .....	41
Table 24:	IVF treatment cycle in which conception occurred, 1979–1997.....	42

Table 25:	Number of oocytes collected by laparoscopy or ultrasound guidance, IVF pregnancies, 1979–1997 .....	42
Table 26:	Number of embryos transferred, IVF pregnancies, 1979–1997 .....	42
Table 27:	Outcome of pregnancy by number of embryos transferred, IVF pregnancies, 1996–1997.....	43
Table 28:	Number of IVF pregnancies following donor oocytes, sperm or embryos, and frozen embryos or oocytes, 1979–1997 .....	43
Table 29:	Outcome of pregnancy after use of donor gametes, donor or frozen embryos, IVF pregnancies, 1979–1997.....	43
Table 30:	Number and outcome of pregnancies after microinsemination, 1990–1997 .....	44
Table 31:	Drugs used in the luteal phase after embryo transfer, IVF pregnancies, 1979–1997.....	44
Table 32:	Outcome of pregnancy by maternal age groups, IVF pregnancies, 1979–1997 ....	45
Table 33:	Spontaneous abortions, IVF pregnancies, 1979–1997 .....	45
Table 34:	Incidence of spontaneous abortions in maternal age groups, IVF pregnancies, 1979–1997 .....	45
Table 35:	Ectopic pregnancies after IVF, 1979–1997.....	46
Table 36:	Heterotopic pregnancies after IVF, 1979–1997.....	46
Table 37:	Reported complications of pregnancy, IVF pregnancies, 1990–1997 .....	46
Table 38:	Duration of singleton and multiple IVF pregnancies of at least 20 weeks' gestation, 1996–1997 .....	47
Table 39:	Duration of singleton and multiple IVF pregnancies of at least 20 weeks' gestation after microinsemination, 1990–1997.....	47
Table 40:	Duration of pregnancy of singleton IVF births after use of donor sperm, donor oocytes or frozen embryos, 1996–1997 .....	48
Table 41:	Maternal age and duration of singleton IVF pregnancies of at least 20 weeks' gestation, 1996–1997 .....	48
Table 42:	Causes of infertility and duration of singleton IVF pregnancies of at least 20 weeks' gestation, 1996–1997 .....	49
Table 43:	Plurality of IVF pregnancies of at least 20 weeks' gestation, 1979–1997.....	49
Table 44:	Plurality of IVF pregnancies of at least 20 weeks' gestation and number of embryos transferred, 1996–1997.....	50
Table 45:	Multiple pregnancies, IVF and GIFT pregnancies, State and Territory, 1996–1997.....	50
Table 46:	Multiple pregnancies in each IVF unit, IVF and GIFT pregnancies, 1996–1997.....	51
Table 47:	Method of delivery for singleton and multiple IVF pregnancies of at least 20 weeks' gestation, 1996–1997 .....	52
Table 48:	Sex of infants in singleton and multiple IVF births of at least 20 weeks' gestation, 1979–1997 .....	52

Table 49:	Birthweight of IVF live births and stillbirths, 1996–1997 .....	52
Table 50:	Birthweight of infants in singleton and multiple IVF births of at least 20 weeks' gestation, 1996–1997 .....	53
Table 51:	Birthweight of infants in singleton and multiple births of at least 20 weeks' gestation after microinsemination, 1990–1997 .....	53
Table 52:	Birthweight of infants in singleton IVF births after use of donor sperm, donor oocytes or frozen embryos, 1979–1997 .....	54
Table 53:	Outcome of infants in singleton and multiple IVF births of at least 20 weeks' gestation, 1996–1997 .....	54
Table 54:	Major congenital malformations in singleton and multiple IVF births of at least 20 weeks' gestation, 1979–1997 .....	54
Table 55:	Numbers and outcomes of GIFT pregnancies by year of conception, 1985–1997.....	55
Table 56:	Place of parental residence, GIFT pregnancies, 1985–1997.....	55
Table 57:	Maternal ages, GIFT pregnancies, 1985–1997 .....	55
Table 58:	Paternal ages, GIFT pregnancies, 1985–1997.....	56
Table 59:	Previous pregnancies for pregnant women, GIFT pregnancies, 1985–1997.....	56
Table 60:	Duration of infertility, GIFT pregnancies, 1985–1997.....	56
Table 61:	Outcome of pregnancy by duration of infertility, GIFT pregnancies, 1996–1997.....	57
Table 62:	Causes of infertility, GIFT pregnancies, 1985–1997 .....	57
Table 63:	Outcome of pregnancy by causes of infertility, GIFT pregnancies, 1996–1997.....	58
Table 64:	Drugs used to stimulate ovulation, GIFT pregnancies, 1985–1997.....	58
Table 65:	GIFT treatment cycle in which conception occurred, 1985–1997.....	58
Table 66:	Number of oocytes collected by laparoscopy or ultrasound guidance, GIFT pregnancies, 1985–1997 .....	59
Table 67:	Number of oocytes transferred, GIFT pregnancies, 1985–1997.....	59
Table 68:	Outcome of pregnancy by number of oocytes transferred, GIFT pregnancies, 1996–1997.....	60
Table 69:	Drugs used in the luteal phase after embryo transfer, GIFT pregnancies, 1985–1997.....	60
Table 70:	Outcome of pregnancy by maternal age groups, GIFT pregnancies, 1985–1997.....	61
Table 71:	Spontaneous abortions, GIFT pregnancies, 1985–1997.....	61
Table 72:	Incidence of spontaneous abortions in maternal age groups, GIFT pregnancies, 1985–1997 .....	61
Table 73:	Ectopic pregnancies after GIFT, 1985–1997 .....	62
Table 74:	Heterotopic pregnancies after GIFT, 1985–1997 .....	62



Table 75:	Reported complications of pregnancy, GIFT pregnancies, 1990–1997 .....	62
Table 76:	Duration of singleton and multiple GIFT pregnancies of at least 20 weeks' gestation, 1996–1997 .....	63
Table 77:	Maternal age and duration of singleton GIFT pregnancies of at least 20 weeks' gestation, 1996–1997 .....	63
Table 78:	Causes of infertility and duration of singleton GIFT pregnancies of at least 20 weeks' gestation, 1996–1997 .....	64
Table 79:	Plurality of GIFT pregnancies of at least 20 weeks' gestation, 1985–1997 .....	64
Table 80:	Plurality of GIFT pregnancies of at least 20 weeks' gestation and number of oocytes transferred, 1996–1997 .....	65
Table 81:	Method of delivery for singleton and multiple GIFT pregnancies of at least 20 weeks' gestation, 1996–1997 .....	65
Table 82:	Sex of infants in singleton and multiple GIFT births of at least 20 weeks' gestation, 1985–1997 .....	65
Table 83:	Birthweight of GIFT live births and stillbirths, 1996–1997 .....	66
Table 84:	Birthweight of infants in singleton and multiple GIFT births of at least 20 weeks' gestation, 1996–1997 .....	66
Table 85:	Outcome of infants in singleton and multiple GIFT births of at least 20 weeks' gestation, 1996–1997 .....	67
Table 86:	Major congenital malformations in singleton and multiple GIFT births of at least 20 weeks' gestation, 1985–1997 .....	67

# Acknowledgments

We thank staff of the IVF units in Australia and New Zealand for completing the data forms and providing additional information. We appreciate the efforts of those individuals, especially clinic co-ordinators and scientists, who have carefully checked records to obtain further details of clinical outcomes, or have provided data on the numbers of women treated, cycles of treatment and laboratory procedures.

We gratefully acknowledge financial support from the Fertility Society of Australia and Serono Australia Pty Ltd. The AIHW National Perinatal Statistics Unit is funded by a grant from the Australian Institute of Health and Welfare to the University of New South Wales.

We thank Professor Michael Chapman, Professor Douglas Saunders, Dr Richard Madden and Dr David Molloy for reviewing the report.

## **Requests for data**

Any enquiries about data for individual IVF units should be directed to the unit concerned. Other enquiries should be made to the AIHW National Perinatal Statistics Unit.

The report may be obtained from:

- Government Info Shops, or AusInfo Mail Order Sales
- Call toll-free on 132 447 or visit <http://www.ausinfo.gov.au/>

# Collaborating IVF and GIFT units

## New South Wales

North Shore Assisted Reproductive Technology, Sydney (Professor Douglas M. Saunders)  
St George Fertility Centre, Sydney (Dr David C. Macourt)  
Lingard Fertility Centre, Newcastle (Dr Robert Woolcott)  
Westmead Fertility Centre, Sydney (Associate Professor Peter Illingworth)  
City West IVF, Sydney (Dr Geoffrey L. Driscoll)  
Royal Prince Alfred Hospital, Sydney (Professor Robert P.S. Jansen)  
Sydney IVF, Sydney (Professor Robert P.S. Jansen)  
Royal Hospital for Women Fertility Group, Sydney (Dr Trevor Johnson)  
Albury Reproductive Medicine Centre, Albury (Dr Scott Giltrap)  
IVF South, Sydney (Professor Michael Chapman)  
Fertility First, Sydney (Dr Anne Clark)

## Victoria

Royal Women's Hospital and Melbourne IVF, Melbourne (Dr Andrew Speirs, Dr Ian Johnston)  
Monash IVF, Melbourne (Professor Gab Kovacs)  
Melbourne Assisted Conception Centre, Mercy Hospital for Women, Melbourne (Dr Mac Talbot)  
Mildura Reproductive Medicine Centre, Mildura (Dr John McBain)

## Queensland

Queensland Fertility Group, Brisbane (Dr David Molloy)  
Monash IVF Gold Coast Fertility Centre, Southport (Dr Irving T. Korman)  
Queensland In-Vitro Fertilization Services, Brisbane (Dr John Allan)  
North Queensland IVF Services, Townsville (Dr Glenn Schaefer)  
Toowoomba IVF, Toowoomba (Dr John Esler)  
IVF Queensland Medical Centre, Buderim (Dr James Dick)

## South Australia

The Queen Elizabeth Hospital, Adelaide (Professor Rob Norman)  
Flinders Reproductive Medicine, Adelaide (Associate Professor Stephen J. Judd)

## Western Australia

PIVET Medical Centre, Perth (Dr John L. Yovich)  
Concept Fertility Centre, Perth (Dr Rob Mazzucchelli)  
Fertility West, Perth (Dr Anne Jequier)

## **Tasmania**

TASIVF, Hobart (Dr Steve Sonneveld)

Launceston SIVF, Launceston (Dr Timothy Sutton)

## **Australian Capital Territory**

Canberra Fertility Centre, Canberra (Dr Martyn A. Stafford-Bell)

## **New Zealand**

Fertility Plus, Auckland (Dr Guy Gudex)

Fertility Associates, Auckland (Dr Richard Fisher, Dr Freddie Graham)

Otago Fertility Services, Dunedin (Associate Professor Wayne Gillett)

ARTEMIS North Shore Fertility, Auckland (Dr Barry Lowe)

The Fertility Centre, Christchurch (Dr Peter Benny)

Fertility Associates, Wellington (Professor John Hutton)

# Abbreviations

NSW	New South Wales
Vic	Victoria
Qld	Queensland
WA	Western Australia
SA	South Australia
Tas	Tasmania
ACT	Australian Capital Territory
NT	Northern Territory
AIHW	Australian Institute of Health and Welfare
NPSU	National Perinatal Statistics Unit
GIFT	gamete intrafallopian transfer
GnRHa	gonadotrophin-releasing hormone analogues
hCG	human chorionic gonadotrophin
ICSI	intracytoplasmic sperm injection
IVF	in-vitro fertilisation
MESA	microepididymal sperm aspiration
na	not available
PESA	percutaneous epididymal sperm aspiration
PROST	pronuclear stage transfer
PZD	partial zona dissection
SUZI	subzonal insemination
TESE	testicular sperm extraction
TEST	tubal embryo stage transfer
ZIFT	zygote intrafallopian transfer



# Highlights

- In Australia, infertile couples were treated by in-vitro fertilisation (IVF) and gamete intrafallopian transfer (GIFT) in 29 units in 1997. New Zealand had 6 IVF units in this period. There were 3,164 births after assisted conception in Australia in 1996, accounting for 1.2% of all births. In New Zealand, there were 254 births after assisted conception in 1996. The number of births continued to increase in 1997, by 9.3% to 3,458 in Australia and by 7.9% to 274 in New Zealand.
- There has been a marked increase in treatment cycles in which intracytoplasmic sperm injection (ICSI) was attempted. Oocyte retrieval cycles for microinsemination increased from 1,243 in 1993 to 2,786 in 1994, 4,261 in 1995, 5,271 in 1996 and 6,308 in 1997. With an additional 3,203 embryo transfer cycles in 1997 after microinsemination and embryo freezing, microinsemination accounted for 40.7% of all transfer cycles in that year. The overall proportion of assisted conception pregnancies resulting from ICSI and other types of microinsemination increased rapidly from less than 1 in 200 (0.4%) in 1990 to 1 in 3 (33.2%) in 1996 and more than 1 in 3 (37.9%) in 1997.
- When all techniques of assisted conception are included together, the viable pregnancy rate in 1997 was 15.2 per 100 embryo transfer cycles, indicating that more than 1 in 7 treatment cycles reaching the stage of embryo transfer resulted in a viable pregnancy of 20 weeks or more.
- In 1997, after transfer of fresh embryos to the uterus, the viable pregnancy rate was 13.3 per 100 oocyte retrieval cycles for IVF without ICSI and 14.7 per 100 oocyte retrieval cycles for ICSI. After GIFT, the viable pregnancy rate was 20.0 per 100 oocyte retrieval cycles.
- After embryo freezing in 1997, the viable pregnancy rate was 12.2 per 100 embryo transfer cycles for IVF without ICSI and 11.4 per 100 embryo transfer cycles for ICSI. After oocyte donation and IVF, the viable pregnancy rate was 15.3 per 100 embryo transfer cycles.
- Between 1996 and 1997, there were slight decreases in the proportion of cycles with more than 3 embryos or oocytes transferred. For all IVF, more than 3 embryos were transferred to the uterus in 2.8% of cycles in 1996 and 2.7% of cycles in 1997; for GIFT, more than 3 oocytes were transferred in 7.2% of cycles in 1996 and 6.9% of cycles in 1997. Women aged 40 years and over accounted for 17.1% of IVF oocyte retrieval cycles, 12.0% of ICSI retrieval cycles, 17.4% of GIFT retrieval cycles, 12.1% of IVF transfer cycles after embryo freezing, 9.1% of ICSI transfer cycles after embryo freezing, and 48.4% of transfer cycles after use of donor oocytes and embryo freezing.
- The multiple pregnancy rate varied markedly between IVF units. In 1996 and 1997, the multiple pregnancy rate for all types of assisted conception ranged from 6.8% to 29.6%. During this same period, multiple pregnancy occurred in 1 in 5 (20.2%) of all IVF and GIFT pregnancies. In IVF pregnancies, twins increased from 17.4% in 1996 to 18.8% in 1997, triplets occurred in 1.4% in both 1996 and 1997, and there were 4 quadruplet pregnancies in 1996. In GIFT pregnancies, twins decreased from 21.0% in 1996 to 18.8% in 1997, triplets decreased from 2.9% in 1996 to 2.5% in 1997, and there was 1 quadruplet pregnancy in 1996.

- In 1996 and 1997, 75 (51.4%) of 146 perinatal deaths after IVF, and 19 (70.4%) of 27 perinatal deaths after GIFT, occurred in multiple births.
- Among 4,476 IVF pregnancies after intracytoplasmic sperm injection and other methods of microinsemination between 1990 and 1997, live births occurred in 77.5%, spontaneous abortion in 19.1%, and ectopic pregnancy in 1.9%. Twins occurred in 17.6% of viable pregnancies, triplets in 1.3%, and there were five quadruplet pregnancies. Among 4,237 ICSI births, the perinatal death rate was 26.2 per 1,000 births; for singleton births, it was 15.8 per 1,000 births; for twins, 42.1 per 1,000 births; and for triplets, 85.1 per 1,000 births. Congenital malformations were reported in 118 (2.8%) fetuses and infants after ICSI, similar to the overall rate among IVF births (2.6%) and GIFT births (2.4%).



# 1 Introduction

This report contains a summary of the results of treatment of infertility by assisted conception in all units in Australia and New Zealand in 1997 and outcomes of pregnancies conceived in 1996 and 1997. The report includes data on in-vitro fertilisation (IVF), gamete intrafallopian transfer (GIFT), and the technique of intracytoplasmic sperm injection (ICSI) used mainly to treat male infertility, but excludes other treatment of infertility by artificial insemination or by ovulation induction without IVF or GIFT. In Australia, there were 29 IVF or GIFT units in 1997. In New Zealand, there were 6 units in 1997. Some IVF units have set up satellite clinics that are linked to major IVF centres in capital cities. Regional centres where satellite clinics have been established include: Albury, Coffs Harbour, Gosford, Lismore, Orange, Wagga Wagga and Wollongong in New South Wales; Ballarat, Bendigo, Broadmeadows, Casterton, Geelong, Maryvale, Morwell, Sale, Shepparton and Wangaratta in Victoria; Cairns, Mackay, Nambour, Rockhampton and Townsville in Queensland; Attadale in Western Australia; and Darwin in the Northern Territory.

The IVF units reported summary data on treatment cycles and also notified each pregnancy on a standard form (Appendix 2). The data included the number of cycles commenced in 1997 and the number progressing to the stages of oocyte retrieval, embryo transfer, clinical pregnancy, and viable pregnancies of at least 20 weeks' gestation. Each IVF unit reported separate results for IVF and uterine transfer of fresh embryos, IVF and tubal transfer of fresh embryos, IVF and transfer of frozen/thawed embryos, donor oocytes, ICSI and GIFT. The tables on treatment cycles are mutually exclusive and so differ slightly from the presentation of IVF results in reports for the years prior to 1991. Each IVF unit was also requested to provide tabulated data on the age distribution, causes of infertility, drugs used to stimulate ovulation, and the number of embryos or oocytes transferred for women treated by IVF, intracytoplasmic sperm injection (ICSI), GIFT and transfer of frozen embryos. Tabulated summaries of results and notified pregnancies were returned to each unit to check their accuracy and completeness.

Each IVF unit is designated by an alphabetical letter which may differ from that given to the unit in previous reports. This code is based on the relative size of the IVF unit (based on the total number of treatment cycles) and therefore may vary from year to year.


All analyses of treatment cycles and pregnancy outcome in this report are based on the year of treatment and conception. Data on pregnancy outcome are given for 1996 and 1997, and include births up to September 1998.

## 1.1 Variations in use of assisted conception

The use of assisted conception to treat infertility can be compared in different populations by relating the number of treatment cycles during a year to the number of women in the reproductive age group. The total number of treatment cycles can be estimated by adding those that reach the stage of oocyte retrieval for IVF and GIFT to the number of transfer cycles for frozen/thawed embryos and donor oocytes. As most women treated by assisted conception are aged between 25 and 44 years, the ratio of the number of treatment cycles is expressed per 100,000 women aged 25-44 years. In the figures for 1997, South Australia and the Northern Territory are reported together because the only IVF clinic in Darwin is a satellite clinic of the Queen Elizabeth Hospital in Adelaide. Also, the figures for Tasmania

and the Australian Capital Territory are combined because there are only three IVF units between the two regions.

There were considerable variations in treatment ratios among the Australian States, and marked differences between Australia and New Zealand (Table 1, Figure 1). In 1997, the treatment ratio in Australia was 835 cycles per 100,000 women, which was 5.2% higher than in 1996. This ratio was nearly three times higher than in New Zealand which had a ratio of 300 per 100,000 women, 7.9% higher than in 1996. In Australia, the highest treatment ratios were in the Australian Capital Territory/Tasmania and Victoria and the lowest ratios were in Queensland and Western Australia. As these ratios are based on the States in which the IVF units are located, comparisons between States may be slightly affected by interstate movements of infertile women for treatment. In Victoria, Tasmania and the Australian Capital Territory, there were relatively more treatment cycles for IVF with fresh embryos than in the other States or New Zealand. Tasmania and the Australian Capital Territory had relatively more IVF with frozen embryos, and Victoria had more treatment cycles using ICSI (fresh or frozen) than in the other States or New Zealand. GIFT was more likely to be used in Queensland than elsewhere.



## 2 Treatment cycles and pregnancy rates

As in previous reports, pregnancy rates are expressed per 100 treatment cycles that reach the stage of oocyte retrieval. In treatment cycles in which embryos were transferred after embryo freezing or oocyte donation, pregnancy rates are expressed per 100 embryo transfer cycles.

Between 1991 and 1996, the total number of treatment cycles for all types of assisted conception increased each year from 16,809 in 1991 to 17,874 in 1992, 18,765 in 1993, 20,706 in 1994, 22,303 in 1995 and 24,121 in 1996. In 1997, there was a further increase of 6.8% to 25,766 cycles.

Following the trend of recent years, the use of microinsemination to treat mainly male infertility continued to increase in 1997. This pattern has been reflected in the resulting viable pregnancies (Figure 2). There were 337 oocyte retrieval cycles for these techniques in 1990, the number then rising each year to 2,786 cycles in 1994, 4,261 cycles in 1995, 5,271 cycles in 1996, and by a further 19.7% to 6,308 cycles in 1997. Microinsemination with transfer of fresh or frozen embryos accounted for 19.7% of all types of assisted conception in 1994, 30.0% in 1995, 35.1% in 1996 and 40.7% in 1997.




Figure 2: Assisted conception pregnancies by method of conception, 1990-1997

The number of transfer cycles after embryo freezing has also continued to increase, from 6,801 cycles in 1996 to 7,723 cycles in 1997, an increase of 13.6%. Transfer cycles after both microinsemination and embryo freezing almost doubled from 929 in 1994 to 1,794 in 1995, increased by a further 26.6% to 2,272 cycles in 1996, and then by another 41.0% to 3,203 cycles in 1997.

## 2.1 IVF and GIFT treatment cycles and pregnancy rates in 1997

Summary data on treatment cycles and pregnancies in 1997 were reported by all IVF units in Australia and New Zealand, but the notifications of pregnancies from one relatively small IVF unit were received too late to be included in this report. The outcomes of all pregnancies conceived in 1996, which were not included in the previous report on assisted conception, are also given in this report. A copy of the pregnancy notification form used in 1996 and 1997 can be found in Appendix 2.

The interpretation of pregnancy rates for the various techniques of assisted conception, and comparison of results in different IVF units, are influenced not only by factors such as age of treated the women and number of embryos or oocytes transferred but also by the relative use of a constantly changing array of techniques. Combining the results for IVF, ICSI and GIFT (but excluding cycles in which frozen embryos or donor oocytes were transferred), the overall viable pregnancy rate was 14.8 per 100 oocyte retrieval cycles (Table 2), slightly higher than the viable pregnancy rate 14.2 per 100 cycles in 1996. When all techniques of assisted conception are included, the viable pregnancy rate for all cycles in which embryos or oocytes were transferred was 15.2 per 100 transfer cycles, compared with 14.4 per 100 transfer cycles in 1996.

IVF units provided summary data on age of the treated women, causes of infertility, drugs used for ovarian stimulation, and number of embryos or oocytes transferred for treatment cycles which progressed to this stage of treatment. Separate data were given for transfers of fresh embryos for IVF, ICSI, and GIFT (Tables 3 and 4) and for frozen embryo transfers for IVF, ICSI and donor oocytes (Table 5). There were relatively more older women among those treated by assisted conception in 1997, continuing the trend of recent years. The proportion of women aged 35 years and over in 1997 was 50.8% for IVF, 44.4% for ICSI and 46.7% for GIFT (Table 3); for women with frozen embryo transfers, the proportion aged 35 years and over was 45.3% for IVF, 38.7% for ICSI and 69.6% for donor oocytes (Table 5). The causes of infertility and the drugs used to stimulate ovulation generally showed a pattern similar to that in previous years. The main causes of infertility were tubal for women treated by IVF, male factor for those treated by ICSI, unexplained infertility for women treated by GIFT and other female causes for those using donor oocytes. For IVF, ICSI and GIFT, the main ovarian stimulants were GnRH analogues, accounting for 92.6%, 92.6% and 93.6%, respectively. In 1997, more than three embryos were transferred in 2.9% of IVF cycles and in 2.3% of ICSI cycles, and more than 3 oocytes were transferred in 6.9% of GIFT cycles (Table 4). More than three thawed embryos were transferred in 0.9% of IVF cycles, in 0.8% of ICSI cycles and in 2.6% of donor oocyte cycles (Table 5).

In 1997, 8,127 treatment cycles were commenced for IVF with a view to subsequent transfer of fresh embryos to the uterus (Table 6). Oocyte retrieval was attempted in 6,697 cycles and embryos were transferred in 5,471 cycles. There were 1,151 clinical pregnancies (17.2 per 100 oocyte retrieval cycles) and 896 viable pregnancies (13.4 per 100 oocyte retrieval cycles). There were marked variations in pregnancy rates among the individual IVF units. This may be partly attributable to differences in selection criteria, methods of treatment, and the characteristics of infertile couples, or to random fluctuations due to the relatively small number of treatment cycles in some IVF units.

An additional 148 treatment cycles were commenced for tubal transfer of embryos after IVF, 12.9% less than the number of cycles commenced in 1996 and less than half the number of cycles commenced in 1995. These resulted in 20 clinical pregnancies (14.1 per 100 oocyte retrieval cycles) and 15 viable pregnancies (10.6 per 100 oocyte retrieval cycles).

In 1997, 33 of the 35 IVF units in Australia and New Zealand used microinsemination techniques to treat infertility. Intracytoplasmic sperm injection (ICSI) was the only type of microinsemination used; no units were using subzonal insemination in 1997. Oocyte retrieval was attempted in 6,234 cycles and embryos were transferred in 5,364 cycles (Table 7), resulting in 1,123 clinical pregnancies (18.0 per 100 oocyte retrieval cycles) and 920 viable pregnancies (14.8 per 100 oocyte retrieval cycles). Pregnancy rates were lower after tubal transfer of ICSI embryos, but there were few treatment cycles in this group (Table 8). Special techniques of sperm collection, such as testicular sperm extraction (TESE), microepididymal sperm aspiration (MESA) and percutaneous epididymal sperm aspiration (PESA), were used in 13.2% of all ICSI treatment cycles (Table 8). Pregnancy rates were generally higher after use of these techniques than for all ICSI cycles.

The 621 embryo transfer cycles after use of donor oocytes (Table 9) resulted in 124 clinical pregnancies (20.0 per 100 embryo transfer cycles) and 95 viable pregnancies (15.3 per 100 embryo transfer cycles).

There were 2,195 treatment cycles commenced for GIFT in 1997 (Table 10), fewer than in previous years, resulting in 506 clinical pregnancies (26.3 per 100 oocyte retrieval cycles) and 384 viable pregnancies (20.0 per 100 oocyte retrieval cycles). There has been relatively little change in GIFT pregnancy rates during the period since the early 1990s in which the use of GIFT has declined.


Data for transfer cycles after embryo freezing are given for conventional IVF (Table 11) and for microinsemination (Table 12). There were 4,520 embryo transfer cycles in the first group and 3,203 cycles in the second group. The total of 7,723 transfer cycles in 1997 continued the increasing trend of previous years (4,607 transfer cycles in 1993, 5,238 in 1994, 6,198 in 1995 and 6,801 in 1996). After conventional IVF and embryo freezing, there were 715 clinical pregnancies (15.8 per 100 embryo transfer cycles) and 551 viable pregnancies (12.2 per 100 embryo transfer cycles). After ICSI and embryo freezing, the clinical and viable pregnancy rates were 14.3 and 11.4 per 100 embryo transfer cycles, respectively.

Embryo freezing avoids the necessity for repeated ovarian stimulation in every treatment cycle. As more couples have their infertility treated by IVF, more embryos are frozen each year. In 1997, more than 6,000 couples decided to have their embryos frozen and more than 32,000 embryos were frozen in that year (Table 13). The number of embryos frozen exceeds the number thawed, thus increasing the total number of embryos in storage each year. In 1997, the average number of embryos frozen was 5.1 per couple, while there were 2.8 embryos per couple transferred after thawing. Some thawed embryos (29.4%) were unsuitable for transfer. By the end of 1997, more than 46,000 frozen embryos were in storage. Policies on how long frozen embryos are kept in storage vary among the IVF units. Comparing changes in the number of stored embryos in different IVF units enables review of these policies.

### 3 IVF pregnancies

This section contains data on all pregnancies other than those resulting from GIFT; it therefore includes pregnancies occurring after transfer of fresh embryos to the uterus or fallopian tubes, transfer of frozen/thawed embryos, use of donor oocytes, and the use of ICSI.

There were 3,038 clinical pregnancies after IVF in 1996 (Table 14, Figure 3), more than in any previous year and an increase of 419 (16.0%) above the number in 1995. There was a further increase of 450 (14.8%) clinical pregnancies to 3,488 in 1997. Live birth was the outcome in 74.2% of the pregnancies in 1996 and in 76.4% in 1997.



### 3.1 Maternal and paternal characteristics


#### 3.1.1 Place of residence

Since 1995, there has been an increase in the number of IVF pregnancies in all Australian States and Territories, except the Northern Territory, and in New Zealand. In 1996 and 1997, nearly 60% of all pregnancies were to couples living in Victoria and New South Wales (Table 15, Figure 4).

#### 3.1.2 Parental age

The majority (80.1%) of women conceiving by IVF in 1996 were in their 30s or 40s (Table 16); this proportion increased slightly to 80.5% in 1997. The proportion of women aged 35 years and over was 40.7% in 1996 and 41.1% in 1997, considerably higher than the proportion of 14.3% for all mothers giving birth in Australia in 1996. Fathers aged 35 years and over increased from 57.9% in 1996 to 59.2% in 1997 (Table 17). The increase in paternal age was

most pronounced for men aged 45 years and over, increasing from 5.9% in previous years to 9.7% in both 1996 and 1997.



### 3.1.3 Previous pregnancies

Women who conceived in 1996 and 1997 had similar previous reproductive experiences to those who conceived in earlier years, but there was a slight increase in the proportion of women who had not been pregnant previously (Table 18).



### 3.1.4 Duration of infertility

There were relatively more women with shorter periods of infertility in 1996 and 1997 than in previous years (Table 19, Figure 5). The proportion of women infertile for a period of less than four years increased from 37.6% in 1979-1995 to 48.4% in 1996 and 52.6% in 1997. Women who had been infertile for 8 years or more were less likely to achieve a live birth than those who had been infertile for shorter periods (Table 20).

### 3.1.5 Causes of infertility

Women conceiving after IVF in 1996 and 1997 were more likely to have infertility due to male factors only or multiple causes than in previous years (19.2% in 1979-1995, 34.3% in 1996, and 33.3% in 1997, and 26.7% in 1979-1995, 30.7% in 1996, and 26.6% in 1997, respectively) and were less likely to have tubal causes (33.6% in 1979-1995, 12.6% in 1996, and 12.3% in 1997) (Table 21, Figure 6). There has been a gradual increase in other stated causes of infertility and unexplained infertility over this period.

The proportion of pregnancies resulting in live births was highest for male infertility (78.9%) and lowest for tubal infertility (72.5%) (Table 22). Spontaneous abortion was highest for endometriosis (22.6%) and lowest for male infertility (17.9%). Ectopic pregnancy was more likely among women treated for tubal causes of infertility (5.5%) than for other causes. Stillbirth was slightly more likely among women treated for tubal infertility (1.2%) than for other causes, but the number of stillbirths in each group was relatively small.



## 3.2 Management of IVF pregnancies

### 3.2.1 Ovarian stimulation

Continuing the trend of recent years, gonadotrophin-releasing hormone analogues (GnRH $\alpha$ ) combined with gonadotrophins were the main drugs used for stimulating ovulation. In 1996 and 1997, these drugs were used in over 90% of treatment cycles that resulted in IVF pregnancies (Table 23, Figure 7). The use of clomiphene to stimulate ovulation has declined



from over 90% of treatment cycles in the mid-1980s to less than 5% of treatment cycles in 1996 and 1997.




Figure 7: Drugs used to stimulate ovulation, IVF pregnancies, 1979–1997

### 3.2.2 Treatment cycle in which pregnancy occurred

Over 40% of all IVF pregnancies occurred in the first treatment cycle (43.5% in 1979–1995, 43.9% in 1996 and 48.9% in 1997), and more than two-thirds of all IVF pregnancies occurred in the first or second treatment cycle (Table 24).

### 3.2.3 Number of oocytes collected

The average number of oocytes collected by laparoscopy or ultrasound guidance for IVF has continued to increase (Table 25, Figure 8). In 1996 and 1997, 15 or more oocytes were collected in over a quarter of all treatment cycles and the mean number of oocytes collected was 11.6 per cycle and 12.0 per cycle, respectively.



### 3.2.4 Number of embryos transferred

There has been a continuing decline in the proportion of IVF pregnancies that resulted from transfer of 3 or more embryos (Figure 9). In 1996, only 2.5% of pregnancies followed transfer of 4 or more embryos; this declined further to 1.7% in 1997 (Table 26). More than 60% of the pregnancies followed transfer of 1 or 2 embryos. The average number of embryos transferred in 1995, 1996 and 1997 was 2.3, less than in previous years (Figure 10).

Of the 138 (2.1%) IVF pregnancies resulting from transfer of 4 or more embryos in 1996 and 1997, 37 (26.8%) resulted in spontaneous abortion, 3 (2.2%) in termination of pregnancy and 2 (1.4%) in stillbirth (Table 27).


### 3.2.5 Donor or frozen gametes and embryos

The number of IVF pregnancies that followed transfer of frozen/thawed embryos has continued to increase from 530 in 1992 to 1,014 in 1996 and 1,192 in 1997 (Table 28). The number of pregnancies after use of donor embryos or donor oocytes increased from 104 in 1995 to 133 in 1997, whereas the number of pregnancies after the use of donor sperm has declined during this period. The outcome of pregnancies after donor sperm, donor oocytes or frozen embryos was similar to that of all IVF pregnancies (Table 29).



### 3.2.6 Microinsemination

IVF pregnancies after microinsemination increased markedly from 1,737 in the six-year period 1990–1995 to 1,224 in 1996 and 1,515 in 1997 (Table 30). There were relatively more live births and fewer ectopic pregnancies after microinsemination than among all IVF pregnancies.



### 3.2.7 Drugs used in luteal phase of pregnancy

Over 80% of women who became pregnant in 1996 and 1997 were treated with drugs during the luteal phase (Table 31). Human chorionic gonadotrophin (hCG) and progestagen were the most commonly used drugs, in 45.0% and 19.1% of IVF pregnancies in 1996 and 44.5% and 20.1% in 1997, respectively.

## 3.3 Outcome of pregnancy

### 3.3.1 Maternal deaths

One woman who conceived by IVF in 1996 died from a pulmonary embolus 8 days after giving birth by caesarean section. There have been three maternal deaths reported in IVF pregnancies among a total of 22,308 pregnancies.

### 3.3.2 Maternal age and outcome of pregnancy

The proportion of women who achieve a live birth after conceiving by IVF decreases with advancing maternal age. Slightly more than three-quarters of women aged less than 35 years gave birth to liveborn infants compared with 74.5% among women aged 35 to 39 years and 59.7% among women aged 40 years and over (Table 32). Spontaneous abortion and


termination of pregnancy were more likely among women of 40 years and over. There was little variation in the occurrence of stillbirths in the various age groups.


### 3.3.3 Spontaneous abortion

There has been little change in the rate of spontaneous abortion in IVF pregnancies in recent years (Table 33, Figure 11). Spontaneous abortion was more likely for older women, increasing from 16.9% for women less than 25 years to 39.7% for women 45 years and over, but only 0.6% of women were in this oldest age group (Table 34, Figure 12). Among 4,476 pregnancies conceived after microinsemination in 1990–1997, there were 853 (19.1%) spontaneous abortions (Table 30).

### 3.3.4 Ectopic pregnancy

The proportion of ectopic pregnancies declined from 4.5% in 1979–1995, to 3.1% in 1996 and 2.6% in 1997 (Table 35, Figure 11). This declining trend is partly attributable to relatively fewer ectopic pregnancies among the increasing proportion of women whose infertility was due to male factors. Among 4,476 pregnancies conceived after microinsemination in 1990–1997, there were 85 (1.9%) ectopic pregnancies (Table 30).





### 3.3.5 Heterotopic pregnancies

Heterotopic pregnancies are those in which there is both a uterine and tubal (ectopic) pregnancy simultaneously. The uterine pregnancy may abort or may continue on to a birth. Heterotopic pregnancies are uncommon. Since assisted conception began in Australia and New Zealand, 169 cases of heterotopic pregnancies (107 leading to abortion and 62 continuing to a birth) have been reported, accounting for 0.5% of all assisted conception pregnancies. There were no reported IVF heterotopic pregnancies prior to 1984 and the first GIFT heterotopic pregnancy was reported in 1986. There have been 14 ICSI heterotopic pregnancies (7 leading to abortion and 7 continuing to a birth). Heterotopic pregnancies after IVF have declined from 0.9% in 1979-1991 to 0.4% in 1996 and 0.2% in 1997 (Table 36).

### 3.3.6 Selective reduction of fetuses

Selective reduction of fetuses may be performed in early pregnancy to abort a severely malformed fetus in a multiple pregnancy or to avoid multiple births. Among pregnancies conceived in 1996, selective reduction was performed in 6 IVF and 2 GIFT pregnancies. Among pregnancies conceived in 1997, there was selective reduction in 9 IVF and 1 GIFT pregnancies. Fetal reduction had previously been performed in 41 pregnancies between 1988 and 1995. Of the 18 pregnancies with selective reduction in 1996-1997, four fetuses were reduced to three in 1 IVF pregnancy and four were reduced to two in 1 IVF and 1 GIFT pregnancies; three fetuses were reduced to two in 8 IVF and 2 GIFT pregnancies and three fetuses were reduced to one in 2 IVF pregnancies; and two fetuses were reduced to one in 3 IVF pregnancies. The indication for fetal reduction was a congenital malformation in 2 IVF pregnancies: trisomy 21 and thalassaemia major, both reduced from two fetuses to one. None of the selective reductions in GIFT pregnancies was for fetal malformations. Among the 18 IVF and GIFT pregnancies in which selective reduction was performed in 1996 and 1997, spontaneous abortion of the remaining fetuses was the outcome in 6 pregnancies.

### 3.3.7 Complications of pregnancy

Significant complications of pregnancy are recorded in tick boxes on the forms used to notify information about the women conceiving by assisted conception and their pregnancies. No

information was given for the data item on complications in 2,131 (12.4%) of the 17,177 pregnancies conceived after IVF in the period from 1990 to 1997. Among the other 15,046 pregnancies for which information was recorded, pregnancy-induced hypertension was reported in 6.2%, threatened abortion in 4.7%, antepartum haemorrhage in 1.8% and placenta praevia in 1.2% (Table 37). Other complications such as maternal medical conditions, fetal growth restriction and premature labour were reported in 15.7% of IVF pregnancies. Any comparison of these reported complications between IVF and other pregnancies should take account of how the information is collected and also the incomplete recording of this data item. Ascertainment of pregnancy complications is improving; this data item was not completed in only 1.9% of pregnancies conceived in 1996–1997 compared with 18.8% in the period from 1990 to 1995.

### **3.3.8 Viable pregnancies of at least 20 weeks' gestation**


Reflecting the overall increase in the number of IVF pregnancies conceived in 1996 and 1997, there was a considerable increase in births. In Australia, there were 3,011 births after IVF in 1997 compared with 2,510 infants conceived in 1996, 2,303 conceived in 1995, and 1,853 conceived in 1994. In New Zealand, the numbers of infants were 267 for conceptions in 1997, 252 for 1996, 169 for 1995, and 154 for 1994.

Preterm births of less than 37 weeks' gestation occurred in 23.8% of all IVF pregnancies in 1996–1997 (Table 38). The incidence of preterm births was higher with increasing plurality, ranging from 13.5% for singleton IVF pregnancies to 62.8% for twin pregnancies and 95.8% for triplet pregnancies. Preterm births among singleton IVF pregnancies declined to their lowest level of 12.7% in 1997 (Figure 13), but this rate was nearly double that for all Australian singleton pregnancies (6.7% in 1996).

After microinsemination, the incidence of preterm births in 1990–1997 was 21.6% for all pregnancies, 12.0% for singleton pregnancies and 59.0% for twin pregnancies (Table 39), slightly less than that for all IVF pregnancies.

In 1996–1997, preterm births occurred in 14.0% of 139 singleton IVF births after use of donor sperm, in 19.9% of 131 births after use of donor oocytes, and in 11.9% of 1,463 births after use of frozen embryos (Table 40).

As in previous years, there was a high proportion of preterm births among singleton IVF pregnancies in all maternal age groups (Table 41) and for all causes of infertility (Table 42). The proportion increased with advancing maternal age from 11.9% for mothers aged 25–29 years to 15.5% for those aged 40 and over. Mothers aged under 25 years had the highest rate of preterm birth (18.8%). Preterm birth was less likely if infertility was due to endometriosis (10.5%) than if it was due to other causes; the highest incidence (14.9%) was among women whose infertility was due to tubal causes.





### 3.3.9 Multiple pregnancies

Multiple pregnancy occurred in 19.0% of IVF pregnancies in 1996 and 20.2% in 1997, similar to the proportion of 19.5% in previous years (Table 43) and much higher than that for all Australian births (1.4% in 1996). The incidence of twin pregnancy after IVF declined from 20.4% in 1988 to 14.4% in 1991, but has since risen again to 17.4% in 1996 and 18.8% in 1997 (Figure 14). The incidence of triplet pregnancy in both 1996 and 1997 was 1.4% (Figure 15). Four quadruplet pregnancies occurred in 1996.


Multiple pregnancy was more likely after transfer of fresh embryos than after transfer of frozen/thawed embryos. In 1996 and 1997, twins occurred in 21.0% of pregnancies after transfer of fresh embryos, triplets in 1.9% and there were 3 quadruplet pregnancies. Among pregnancies after transfer of frozen embryos, twins occurred in 12.7%, triplets in 0.4% and there was 1 quadruplet pregnancy.

The likelihood of multiple pregnancy depends on the number of embryos transferred. In 1996 and 1997, twins occurred in 18.7% of IVF pregnancies after transfer of 2 embryos, in 21.3% after transfer of 3 embryos, and in 18.5% after transfer of 4 embryos (Table 44). Among 3,511 pregnancies conceived after microinsemination in 1990–1997, there were 669 multiple pregnancies (19.1%); twins occurred in 617 (17.6%), triplets in 47 (1.3%) and quadruplets in 5 (0.1%).



Multiple pregnancies after assisted conception were slightly less common in Australia (19.9%) than in New Zealand (23.0%) (Table 45, Figure 16). Queensland and Western Australia had the highest multiple pregnancy rates, 23.3% and 22.1%, respectively, and New South Wales had the lowest, 18.0%. There were relatively more triplets and quadruplets in Queensland (3.1%) and Western Australia (2.4%) than in other States.






Multiple pregnancy rates varied considerably among the IVF units, ranging from 6.8% to 29.6%, with one small unit having no multiple pregnancies (Table 46). Some of this variability may be due to the relatively small numbers of pregnancies reported in many of the units (57.1% reported fewer than 100 pregnancies, and 74.3% reported fewer than 200 in the years 1996 and 1997). When only the larger IVF units with at least 100 reported pregnancies were considered, multiple pregnancy rates ranged from 13.1% (among 145 pregnancies) to 29.6% (among 165 pregnancies). Overall, multiple pregnancies occurred in 20.3% of the total of 5,870 IVF and GIFT pregnancies in the two-year period.

### 3.3.10 Method of delivery

As in previous years, caesarean rates were higher for multiple than for singleton IVF pregnancies (Figure 17). In 1996–1997, the caesarean rate was 35.8% for singleton pregnancies, 61.0% for twin pregnancies and 87.3% for triplet pregnancies (Table 47). The caesarean rate for singleton IVF pregnancies was considerably higher than the rate of 19.1% for singleton Australian births in 1996. The caesarean rate for singleton IVF pregnancies increased with maternal age, from 29.8% for mothers under 30 years of age to 54.8% for mothers aged 40 years and over.

### 3.3.11 Sex of infants

The sex ratio of infants born after IVF was 105.7 in 1996 and 108.7 in 1997, similar to previous years (Table 48). The sex ratio of infants born in all years after use of donor sperm was 101.4 among 1,444 births; after use of donor oocytes, it was 115.6 among 565 births; and after use of frozen embryos, it was 108.1 among 5,269 births. Among the 4,237 births after microinsemination, the sex ratio was 95.1.



### 3.3.12 Birthweight

The mean birthweight and the incidence of low birthweight (less than 2500 g) for infants born after IVF in 1996 and 1997 differed considerably from the birthweights for all Australian births in 1996. The mean birthweight of IVF births in 1996 and 1997 was 2,900 g (Table 49), 460 g less than the mean birthweight of 3,360 g for all Australian births in 1996. The high incidence of multiple births after IVF accounted for much of this difference (Table 50). For singleton births, the mean birthweight was 3,236 g after IVF and 3,389 g for all Australian births; for twins, 2,313 g and 2,392 g, respectively; and for triplets, 1,700 g and 1,642 g, respectively. Among singleton IVF births in 1996 and 1997, low birthweight occurred in 10.6%, compared with 5.1% for all singleton births in Australia in 1996.

The mean birthweight of 2,842 singleton births after microinsemination was 3,248 g (Table 51), similar to that for all singleton IVF births. Low birthweight occurred in 9.9% of singleton births after microinsemination (Table 51), in 10.5% after donor sperm, in 13.2% after donor oocytes, and in 7.6% after embryo freezing (Table 52).

There were relatively fewer low birthweight infants born after IVF than after GIFT (Figure 18). The incidence of low birthweight in singleton IVF births was lowest in 1994, then slightly higher in subsequent years (Figure 19).

### 3.3.13 Perinatal mortality

Perinatal deaths include fetal deaths (stillbirths) of at least 20 weeks' gestation and neonatal deaths of liveborn infants occurring within 28 days of birth. The perinatal death rate for all IVF births in 1996–1997 was 24.2 per 1,000 births (Table 53), lower than in previous years (Figure 20); for singleton births, the rate was 17.7 per 1,000 births, for twins, 34.3 per 1,000 births, and for triplets, 51.6 per 1,000 births. Among 4,237 births after microinsemination in 1990–1997, the perinatal death rate was 26.2 per 1,000 births; for singleton births, the rate was 15.8 per 1,000 births, for twins, 42.1 per 1,000 births, and for triplets, 85.1 per 1,000 births.

The perinatal death rate after assisted conception is considerably higher than for all births. In 1997, the perinatal death rate among all births of at least 20 weeks' gestation or 400 g

birthweight in Australia was 9.2 per 1,000 births (Australian Bureau of Statistics 1999). Factors contributing to the higher perinatal death rate after assisted conception include relatively more older mothers, their underlying causes of infertility, and the much higher incidence of multiple births than in the general population.




Figure 18: Birthweight of IVF and GIFT births, 1979-1997




Figure 19: Incidence of low birthweight in singleton IVF births, 1979-1997




Figure 20: Perinatal mortality in singleton and multiple IVF births, 1979–1997


### 3.3.14 Congenital malformations

Among 20,376 live births, stillbirths and induced abortions of at least 16 weeks' gestation after IVF in 1979–1997, there were 529 (2.6%) infants and fetuses with major congenital malformations (Table 54). The malformation rate was higher in singleton births (3.0%) than in multiple births (1.7%).

Among pregnancies conceived in 1990–1997 after microinsemination, there were 18 pregnancies terminated for fetal abnormality and 4,237 births. Major congenital malformations were notified in 118 fetuses and infants, a malformation rate of 2.8%. There were 98 (3.4%) malformations among 2,860 singleton births and 20 (1.4%) among 1,395 multiple births.

## 4 GIFT pregnancies

The number of GIFT pregnancies has declined substantially since 1994, after showing little change between 1991 and 1993 (Table 55, Figure 21). There were 505 GIFT pregnancies in 1997, about half the number conceived in 1991, 1992 and 1993. In 1996, 78.3% of GIFT pregnancies resulted in live births; in 1997, there were relatively more early pregnancy losses and live births occurred in 71.5% of pregnancies.



### 4.1 Maternal and paternal characteristics

#### 4.1.1 Place of residence

In 1996 and 1997, there were relatively more GIFT pregnancies in Queensland, increasing from 28.5% in 1985–1995 to 36.4% in 1996 and 43.8% in 1997 (Table 56). There was a decline in the proportion in most States; 75% of all GIFT pregnancies occurred in Queensland or New South Wales.

#### 4.1.2 Parental age

Women who conceived after GIFT in 1996 and 1997 were older than those in previous years, increasing from 30.0% aged 35 years and over in 1985–1995 to 37.8% in 1996 and 41.2% in 1997 (Table 57). Paternal age also increased from 46.9% aged 35 years and over in 1985–1995 to 51.8% in 1996 and 55.0% in 1997 (Table 58). Fathers were relatively older than mothers, with more than three times as many fathers aged 40 years and over. In 1985–1997, 5.7% of mothers and 18.7% of fathers were in this age group. Women who became pregnant after GIFT were generally older than the mothers of all babies born in Australia, and slightly younger than those who became pregnant after IVF. In 1996 and 1997, 39.3% of GIFT

pregnancies were to women aged 35 years and over, compared with 40.9% of IVF pregnancies and 14.3% of all Australian births.

### 4.1.3 Previous pregnancies

Women who conceived by GIFT in recent years were slightly more likely to have been pregnant more than once previously than those who had conceived by GIFT in earlier years (Table 59).

### 4.1.4 Duration and causes of infertility

Among women conceiving by GIFT in 1996 and 1997, more than 50% had been infertile for periods of less than 4 years, relatively more than in previous years (Table 60). The likelihood of spontaneous abortion was highest among those with a duration of infertility of 8 or more years; these women were also less likely to have a live birth (69.7%) than those who had a shorter duration of infertility (Table 61).

The causes of infertility that preceded GIFT pregnancies in 1996 and 1997 varied slightly from those in previous years (Table 62, Figure 22). The proportion of GIFT pregnancies resulting in live births was similar for all causes of infertility in 1996 and 1997 (Table 63). There was a low incidence of ectopic pregnancy among women whose infertility was due to endometriosis (0.6%) or male factor (0.5%). The highest incidence of ectopic pregnancy after GIFT was associated with tubal causes of infertility (6.9%).




Figure 22: Causes of infertility, GIFT pregnancies, 1985-1997

## 4.2 Management of GIFT pregnancies

### 4.2.1 Ovarian stimulation

The trend of increasing use of GnRH analogues (97.2% in 1996 and 94.5% in 1997) and declining use of clomiphene (1.4% and 2.0%, respectively) to induce ovulation (Table 64) was

similar to that for IVF pregnancies (Figure 7). Over half (52.2%) of GIFT pregnancies occurred in the first treatment cycle in 1996 and 1997, and another quarter (23.6%) occurred in the second cycle, slightly higher than in previous years (Table 65).

#### 4.2.2 Number of oocytes collected and transferred

The mean number of oocytes collected in treatment cycles that resulted in GIFT pregnancies was 8.4 in 1996 and 8.7 in 1997 (Table 66), higher than in previous years but much lower than the means of 11.6 and 12.0, respectively, for IVF pregnancies (Figure 8). In 1996 and 1997, 15 or more oocytes were collected in 10.8% of all treatment cycles, higher than in previous years (7.7%).

In 1997, 93.4% of GIFT pregnancies followed transfer of three or fewer oocytes (Table 67), similar to 1996 (93.3%) and much higher than in previous years (77.3%). In the 1990s, there has been an increasing trend in pregnancies occurring after transfer of 2 oocytes, increasing from 10.6% in 1990 to 45.1% in 1997 (Figure 23). The proportion of pregnancies with live births varied with the number of oocytes transferred (Table 68). Women having pregnancies following the transfer of 1 oocyte or at least 4 oocytes were the least likely to have a live birth (54.5% and 66.2%, respectively), while women having 2 or 3 oocytes transferred were the most likely to have a live birth (76.3% and 77.1%, respectively).




Figure 23: Percentage distribution of number of oocytes transferred, GIFT pregnancies, 1985-1997

#### 4.2.3 Drugs used in luteal phase of pregnancy

Most women (98.2%) who became pregnant after GIFT in 1996 and 1997 were treated with drugs during the luteal phase (Table 69). Human chorionic gonadotrophin (hCG) and Proluton were the most commonly used drugs.

## **4.3 Outcome of pregnancy**

### **4.3.1 Maternal deaths**

Three maternal deaths have been reported in GIFT pregnancies, but there were no deaths in 1996 or 1997.

### **4.3.2 Spontaneous abortion and ectopic pregnancy**

Among almost 8,800 GIFT pregnancies conceived between 1985 and 1997, spontaneous abortion was the outcome in 20.7% (Table 70).

In 1996, spontaneous abortion occurred in 19.8% of intrauterine GIFT pregnancies and this increased to 23.9% in 1997, higher than in previous years (Table 71). Older women had higher rates of spontaneous abortion, 44.4% for those aged 40–44 years and 60.0% for those aged 45 and over (Table 72).

Ectopic pregnancy occurred in 1.7% of GIFT pregnancies in 1996, fewer than in previous years (Table 73). In 1997, the incidence of ectopic pregnancies after GIFT increased to 3.2%.

### **4.3.3 Heterotopic pregnancies**

Heterotopic pregnancies after GIFT declined from 0.8% in 1985–1991 to 0.2% in 1994 and 1995, but then increased to 0.5% in 1996 and 0.8% in 1997 (Table 74).

### **4.3.4 Complications of pregnancy**

The comments under this heading in Section 3.3.7 should be noted. Information was not recorded for this data item in 1,176 (18.0%) of the 5,359 pregnancies conceived after GIFT between 1990 and 1997. Ascertainment of pregnancy complications is improving, only 2.4% of notifications of GIFT pregnancies lacking completion of this data item in 1996–1997. Among 5,348 GIFT pregnancies for which information was recorded, pregnancy-induced hypertension was reported in 6.6%, threatened abortion in 6.0%, antepartum haemorrhage in 2.4%, placenta praevia in 0.9% and other complications in 19.0% (Table 75).

### **4.3.5 Viable pregnancies of at least 20 weeks' gestation**

Reflecting the overall decline in the number of GIFT pregnancies in recent years, the number of births in Australia after GIFT decreased from 1,023 in 1992 to 654 in 1996 and 447 in 1997. There were very few births after GIFT in New Zealand, from 9 in 1992, to 5 in 1993, 7 in 1994, 6 in 1995, 2 in 1996 and 7 in 1997. The total of 3,164 births after IVF and GIFT in Australia in 1996 represented 1.2% of the national births.

Preterm birth occurred in 23.2% of all GIFT pregnancies in 1996 and 1997 (Table 76), similar to other years, and in 12.5% of singleton pregnancies, in 53.1% of twin pregnancies, and in 100.0% of triplet pregnancies. In singleton GIFT pregnancies, there was a lower incidence of preterm birth for births to mothers aged 25–29 years (8.3%), while the incidence of preterm birth was higher in the older maternal age groups (Table 77). Relatively lower rates of preterm birth occurred when infertility was due to male factors (4.3%); the incidence of preterm birth was similar for all other causes of infertility (Table 78).



### **4.3.6 Multiple pregnancies**

Twins occurred in 21.0% of GIFT pregnancies in 1996 and in 18.8% in 1997 (Table 79), slightly lower than the rate in recent years (Figure 14). The incidence of triplets after GIFT was 2.9% in 1996 and 2.5% in 1997, still higher than in IVF pregnancies (Figure 15). There was also 1 quadruplet pregnancy in 1996.

In 1996 and 1997, as in previous years, higher multiple pregnancy rates were usually associated with transfer of more oocytes (Table 80). There were 2 (16.7%) twin pregnancies after transfer of one oocyte, 16.8% after transfer of two oocytes, 24.2% after three oocytes, and 14.0% after transfer of 4 oocytes.

### **4.3.7 Method of delivery**

Caesarean birth rates for GIFT pregnancies were higher for multiple births than for singleton births (Table 81). In 1996 and 1997, the caesarean rate was 33.3% in singleton pregnancies, 66.1% in twin pregnancies, and 79.2% in triplet pregnancies. In singleton GIFT pregnancies, the caesarean rate was 19.7% for mothers aged less than 25 years, 26.3% for those aged 25–29, 30.6% for those aged 30–34, 41.1% for those aged 35–39, and 47.4% for mothers aged 40 years and over.

### **4.3.8 Sex of infants**

The sex ratio of infants born after GIFT in 1996 was 119.4, higher than in previous years (Table 82). In 1997, the sex ratio was 109.3, decreasing from 112.6 for singletons to 109.1 for twins and 80.0 for triplets. In recent years, the sex ratio of infants born after GIFT has been consistently higher than the sex ratio for IVF.

### **4.3.9 Infant's birthweight**

The mean birthweight of 2,824 g after GIFT in 1996 and 1997 (Table 83) was slightly higher than that of 2,803 g in the previous year but less than the average of 2,900 g after IVF. The high incidence of multiple births after GIFT accounted for much of this difference (Table 84). In 1996 and 1997, for singleton GIFT births, the mean birthweight was 3,209 g; for twins, it was 2,367 g; and for triplets, it was 1,532 g.


In 1996 and 1997, low birthweight occurred in 29.5% of all GIFT pregnancies (Table 84), slightly less than the rate in 1995 (31.3%). There were relatively more low birthweight infants born after GIFT than after IVF (Figure 18). Low birthweight was more likely with increasing plurality, ranging from 9.6% for singleton GIFT births to 52.7% for twins and 98.6% for triplets.

### **4.3.10 Perinatal mortality**

The perinatal death rate after GIFT in 1996 and 1997 was 24.3 per 1,000 births (Table 85), lower than the total rate of 29.5 per 1,000 births in 1995. There were relatively fewer stillbirths; the stillbirth rate was 15.3 per 1,000 births compared with 20.2 per 1,000 in 1995 (Figure 24). Of the 27 perinatal deaths after GIFT in 1996 and 1997, 19 (70.4%) were in multiple births.

### 4.3.11 Congenital malformations

Major congenital malformations occurred in 209 (2.4%) of 8,608 births and induced abortions after GIFT in the period from 1985 to 1997 (Table 86). The malformation rate of 2.7% in singleton births was slightly higher than that of 2.0% in multiple births.



## 5 Tables

**Table 1: Use of assisted conception to treat infertility, selected States, Australia and New Zealand, 1997**

State / Country	Women aged 25–44 years (thousands)	Treatment cycles <sup>(a)</sup>	Ratios <sup>(b)</sup>
New South Wales	967	7,986	826
Victoria	723	7,349	1,017
Queensland	523	3,399	649
Western Australia	283	1,880	665
South Australia <sup>(c)</sup>	255	2,127	834
Australian Capital Territory and Tasmania	122	1,252	1,023
<b>Australia</b>	<b>2,874</b>	<b>23,993</b>	<b>835</b>
<b>New Zealand</b>	<b>592</b>	<b>1,773</b>	<b>300</b>

(a) Includes IVF fresh, IVF frozen, GIFT, ICSI, Tubal transfer and Donor oocytes/embryos

(b) Treatment cycles per 100,000 women aged 25–44 years

(c) Includes external unit based in the Northern Territory

**Table 2: Viable pregnancy rates for all techniques of assisted conception, 1997**

Type of assisted conception	Oocyte retrieval cycles	Embryo/ gamete transfer cycles	Viable pregnancies	Viable pregnancy rate per 100 oocyte retrievals	Viable pregnancy rate per 100 embryo transfers
IVF: fresh embryos	6,839	5,593	911	13.3	16.3
ICSI: fresh embryos	6,308	5,438	928	14.7	17.1
IVF: frozen embryos	-	4,520	551	-	12.2
ICSI: frozen embryos	-	3,203	364	-	11.4
IVF: donor oocytes	-	621	95	-	15.3
GIFT	1,924	1,858	384	20.0	20.7
<b>All techniques</b>	<b>15,071 <sup>(a)</sup></b>	<b>21,233</b>	<b>3,233</b>	<b>14.8 <sup>(a)</sup></b>	<b>15.2</b>

(a) Exclude data on cycles with frozen embryos and donor oocytes

**Table 3: Oocyte retrieval cycles for IVF, ICSI and GIFT, by maternal age, cause of infertility, and drugs used to stimulate ovulation, 1997**

Characteristic	Oocyte retrieval cycles attempted					
	IVF		ICSI		GIFT	
	Number	Per cent	Number	Per cent	Number	Per cent
<b>Maternal age (at start of treatment)</b>						
<20	1	0.0	5	0.1	-	-
20-24	82	1.2	168	2.7	31	1.6
25-29	915	13.8	1,158	18.8	326	17.0
30-34	2,267	34.1	2,098	34.0	665	34.7
35-39	2,240	33.7	1,996	32.4	562	29.3
40-44	1,046	15.7	680	11.0	315	16.4
45+	93	1.4	63	1.0	20	1.0
Not stated	4		-		-	
<b>All ages</b>	<b>6,648</b>	<b>100.0</b>	<b>6,168</b>	<b>100.0</b>	<b>1,919</b>	<b>100.0</b>
<b>Cause(s) of infertility</b>						
Tubal only	2,120	31.9	226	3.6	65	3.4
Other female only	924	13.9	241	3.8	463	24.1
Male factors only	839	12.6	3,811	60.3	260	13.5
Multiple causes	1,227	18.4	1,405	22.2	279	14.5
Unexplained	1,546	23.2	642	10.2	852	44.4
Not stated	3		-		-	
<b>All causes</b>	<b>6,659</b>	<b>100.0</b>	<b>6,325</b>	<b>100.0</b>	<b>1,919</b>	<b>100.0</b>
<b>Ovarian stimulation</b>						
GnRH analogues + other	6,154	92.6	5,858	92.6	1,793	93.5
No GnRH analogues						
— clomiphene + any other	289	4.3	370	5.8	66	3.4
— other drugs	27	0.4	16	0.3	14	0.7
— natural cycles	176	2.6	81	1.3	44	2.3
Not stated	13		-		2	
<b>Total</b>	<b>6,659</b>	<b>100.0</b>	<b>6,325</b>	<b>100.0</b>	<b>1,919</b>	<b>100.0</b>

**Table 4: Embryo transfer cycles for IVF, ICSI and GIFT, by number of embryos or oocytes transferred, 1997**

Number of embryos / oocytes transferred	Embryo transfer cycles					
	IVF		ICSI		GIFT	
	Number	Per cent	Number	Per cent	Number	Per cent
One	798	14.3	720	13.2	101	5.5
Two	2,828	50.6	2,924	53.7	855	46.1
Three	1,799	32.2	1,679	30.8	769	41.5
Four	159	2.8	119	2.2	106	5.7
Five	7	0.1	6	0.1	4	0.2
Six or more	2	0.0	2	0.0	18	1.0
Not stated	4		-		-	
<b>Total</b>	<b>5,597</b>	<b>100.0</b>	<b>5,450</b>	<b>100.0</b>	<b>1,853</b>	<b>100.0</b>

**Table 5: Number of transfer cycles after cryopreservation, by maternal age, cause of infertility, and number of embryos transferred, 1997**

Characteristic	Embryo transfer cycles					
	IVF		ICSI		Donor oocytes	
	Number	Per cent	Number	Per cent	Number	Per cent
<b>Maternal age (at start of treatment)</b>						
<20	4	0.1	1	0.0	-	-
20-24	62	1.4	68	2.4	1	0.3
25-29	657	14.9	613	21.5	23	6.7
30-34	1,689	38.3	1,069	37.4	81	23.5
35-39	1,465	33.2	843	29.5	73	21.2
40-44	505	11.4	247	8.7	115	33.3
45+	29	0.7	14	0.5	52	15.1
Not stated	276		170		10	
<b>All ages</b>	<b>4,687</b>	<b>100.0</b>	<b>3,025</b>	<b>100.0</b>	<b>355</b>	<b>100.0</b>
<b>Cause(s) of infertility</b>						
Tubal only	1,133	26.9	53	2.0	15	4.7
Other female only	637	15.1	59	2.3	129	40.1
Male factors only	577	13.7	1,670	64.1	32	9.9
Multiple causes	751	17.9	639	24.5	89	27.6
Unexplained	1,108	26.3	185	7.1	57	17.7
Not stated	481		419		33	
<b>All causes</b>	<b>4,687</b>	<b>100.0</b>	<b>3,025</b>	<b>100.0</b>	<b>355</b>	<b>100.0</b>
<b>Number of embryos transferred</b>						
One	930	20.9	624	21.8	90	25.6
Two	2,593	58.4	1,741	60.7	183	52.0
Three	877	19.7	478	16.7	70	19.9
Four	42	0.9	22	0.8	9	2.6
Five	1	0.0	1	0.0	-	-
Six or more	-	-	-	-	-	-
Not stated	244		159		3	
<b>Total</b>	<b>4,687</b>	<b>100.0</b>	<b>3,025</b>	<b>100.0</b>	<b>355</b>	<b>100.0</b>

**Table 6: IVF pregnancies after transfer of fresh embryos to uterus, numbers and pregnancy rates in each IVF unit, 1997**

Stage of treatment	IVF unit												
	A	B	C	D	E	F	G	H	I	J	K	L	M
Treatment cycles commenced	1,196	1,258	582	507	247	363	432	296	289	144	275	222	147
Cycles with oocyte retrieval	908	1,024	493	444	225	333	338	219	233	133	221	201	130
Cycles with embryo transfer	714	749	426	414	174	263	294	191	192	86	194	163	112
Clinical pregnancies	119	157	80	100	30	97	69	38	27	10	42	22	18
Viable pregnancies	96	121	40	79	25	75	61	31	24	6	37	17	14
Clinical pregnancies per 100 oocyte retrieval cycles	13.1	15.3	16.2	22.5	13.3	29.1	20.4	17.4	11.6	7.5	19.0	10.9	13.8
Viable pregnancies per 100 oocyte retrieval cycles	10.6	11.8	8.1	17.8	11.1	22.5	18.0	14.2	10.3	4.5	16.7	8.5	10.8

Stage of treatment	IVF unit												
	N	O	P	Q	R	S	T	U	V	W	X	Y	Z
Treatment cycles commenced	287	58	198	60	190	216	82	80	155	84	112	104	56
Cycles with oocyte retrieval	212	52	166	54	183	161	75	65	138	82	90	87	50
Cycles with embryo transfer	168	48	129	43	165	141	61	49	117	77	73	69	44
Clinical pregnancies	49	14	33	7	49	28	6	16	28	20	16	13	4
Viable pregnancies	41	10	26	6	34	26	5	9	22	18	13	11	2
Clinical pregnancies per 100 oocyte retrieval cycles	23.1	26.9	19.9	13.0	26.8	17.4	8.0	24.6	20.3	24.4	17.8	14.9	8.0
Viable pregnancies per 100 oocyte retrieval cycles	19.3	19.2	15.7	11.1	18.6	16.1	6.7	13.8	15.9	22.0	14.4	12.6	4.0

Stage of treatment	IVF unit										Total
	AA	AB	AC	AD	AE	AF	AG	AH	AI		
Treatment cycles commenced	98	-	114	53	40	56	30	40	56		8,127
Cycles with oocyte retrieval	65	-	89	48	37	54	15	29	43		6,697
Cycles with embryo transfer	57	-	72	43	25	48	14	24	32		5,471
Clinical pregnancies	15	-	12	13	3	6	-	7	3		1,151
Viable pregnancies	11	-	12	8	3	5	-	5	3		896
Clinical pregnancies per 100 oocyte retrieval cycles	23.1	-	13.5	27.1	8.1	11.1	-	24.1	7.0		17.2
Viable pregnancies per 100 oocyte retrieval cycles	16.9	-	13.5	16.7	8.1	9.3	-	17.2	7.0		13.4

**Table 7: IVF pregnancies after ICSI, numbers and pregnancy rates in each IVF unit, 1997**

Stage of treatment	IVF unit												
	A	B	C	D	E	F	G	H	I	J	K	L	M
Treatment cycles commenced	982	991	378	576	350	329	163	263	151	197	293	199	147
Cycles with oocyte retrieval	982	870	378	530	341	308	145	219	137	188	226	178	147
Cycles with embryo transfer	825	694	360	490	286	261	129	171	108	154	198	167	130
Clinical pregnancies	140	155	96	121	54	83	27	28	22	25	53	29	21
Viable pregnancies	113	129	77	108	39	68	24	19	19	16	46	21	13
Clinical pregnancies per 100 oocyte retrievals	14.3	17.8	25.4	22.8	15.8	26.9	18.6	12.8	16.1	13.3	23.5	16.3	14.3
Viable pregnancies per 100 oocyte retrievals	11.5	14.8	20.4	20.4	11.4	22.1	16.6	8.7	13.9	8.5	20.4	11.8	8.8

Stage of treatment	IVF unit												
	N	O	P	Q	R	S	T	U	V	W	X	Y	Z
Treatment cycles commenced	86	240	107	235	103	68	112	116	73	129	69	78	64
Cycles with oocyte retrieval	86	234	86	211	98	61	111	97	64	128	53	62	58
Cycles with embryo transfer	79	207	69	171	87	50	105	80	60	120	47	51	52
Clinical pregnancies	20	29	10	34	25	9	20	12	15	25	9	10	11
Viable pregnancies	19	23	8	27	21	8	18	11	13	23	6	8	8
Clinical pregnancies per 100 oocyte retrievals	23.3	12.4	11.6	16.1	25.5	14.8	18.0	12.4	23.4	19.5	17.0	16.1	19.0
Viable pregnancies per 100 oocyte retrievals	22.1	9.8	9.3	12.8	21.4	13.1	16.2	11.3	20.3	18.0	11.3	12.9	13.8

Stage of treatment	IVF unit										Total
	AA	AB	AC	AD	AE	AF	AG	AH	AI		
Treatment cycles commenced	-	-	28	52	46	27	46	47	8		6,753
Cycles with oocyte retrieval	-	-	28	49	44	26	36	45	8		6,234
Cycles with embryo transfer	-	-	23	49	31	26	32	44	8		5,364
Clinical pregnancies	-	-	4	8	4	7	6	10	1		1,123
Viable pregnancies	-	-	4	8	4	7	4	7	1		920
Clinical pregnancies per 100 oocyte retrievals	-	-	14.3	16.3	9.1	26.9	16.7	22.2	12.5		18.0
Viable pregnancies per 100 oocyte retrievals	-	-	14.3	16.3	9.1	26.9	11.1	15.6	12.5		14.8

**Table 8: Pregnancy rates by type of transfer and technique of sperm collection, ICSI pregnancies, 1997**

Stage of treatment	Type of transfer		Sperm collection technique			
	Uterine	Tubal	TESE	MESA	PESA	Other
Treatment cycles commenced	6,753	87	490	290	116	6
Cycles with oocyte retrieval	6,234	74	484	269	115	6
Cycles with embryo transfer	5,364	74	412	247	104	6
Clinical pregnancies	1,123	12	83	63	30	2
Viable pregnancies	920	8	74	54	22	1
Clinical pregnancies per 100 oocyte retrievals	18.0	16.2	17.1	23.4	26.1	33.3
Viable pregnancies per 100 oocyte retrievals	14.8	10.8	15.3	20.1	19.1	16.7

**Table 9: IVF pregnancies after use of donor oocytes, numbers and pregnancy rates in each IVF unit, 1997**

Stage of treatment	IVF unit												
	A	B	C	D	E	F	G	H	I	J	K	L	M
Cycles with embryo transfer	138	141	52	28	3	7	38	23	6	31	11	8	25
Clinical pregnancies	23	33	7	7	1	4	6	4	1	6	4	1	7
Viable pregnancies	21	23	5	3	1	4	6	2	1	4	4	1	3
Clinical pregnancies per 100 embryo transfers	16.7	23.4	13.5	25.0	33.3	57.1	15.8	17.4	16.7	19.4	36.4	12.5	28.0
Viable pregnancies per 100 embryo transfers	15.2	16.3	9.6	10.7	33.3	57.1	15.8	8.7	16.7	12.9	36.4	12.5	12.0

Stage of treatment	IVF unit												
	N	O	P	Q	R	S	T	U	V	W	X	Y	Z
Cycles with embryo transfer	1	14	6	11	10	16	-	4	6	3	10	-	4
Clinical pregnancies	-	1	-	3	4	2	-	-	-	-	2	-	1
Viable pregnancies	-	1	-	2	4	2	-	-	-	-	-	-	1
Clinical pregnancies per 100 embryo transfers	-	7.1	-	27.3	40.0	12.5	-	-	-	-	20.0	-	25.0
Viable pregnancies per 100 embryo transfers	-	7.1	-	18.2	40.0	12.5	-	-	-	-	-	-	25.0

Stage of treatment	IVF unit										Total
	AA	AB	AC	AD	AE	AF	AG	AH	AI		
Cycles with embryo transfer	9	-	2	-	7	-	6	-	1		621
Clinical pregnancies	3	-	-	-	1	-	2	-	1		124
Viable pregnancies	3	-	-	-	1	-	2	-	1		95
Clinical pregnancies per 100 embryo transfers	33.3	-	-	-	14.3	-	33.3	-	100.0		20.0
Viable pregnancies per 100 embryo transfers	33.3	-	-	-	14.3	-	33.3	-	100.0		15.3



**Table 10: GIFT pregnancies, numbers and pregnancy rates in each IVF unit, 1997**

Stage of treatment	IVF unit												
	A	B	C	D	E	F	G	H	I	J	K	L	M
Treatment cycles commenced	21	191	607	86	235	189	90	1	2	148	16	126	5
Cycles with oocyte retrieval	21	156	537	80	205	172	69	1	2	142	14	101	5
Cycles with gamete transfer	21	146	528	80	182	170	67	1	2	136	14	98	5
Clinical pregnancies	2	34	176	23	48	45	6	1	-	36	3	19	2
Viable pregnancies	2	30	127	16	37	39	5	1	-	25	3	15	1
Clinical pregnancies per 100 oocyte retrieval cycles	9.5	21.8	32.8	28.8	23.4	26.2	8.7	100.0	-	25.4	21.4	18.8	40.0
Viable pregnancies per 100 oocyte retrieval cycles	9.5	19.2	23.6	20.0	18.0	22.7	7.2	100.0	-	17.6	21.4	14.9	20.0

Stage of treatment	IVF unit												
	N	O	P	Q	R	S	T	U	V	W	X	Y	Z
Treatment cycles commenced	-	38	22	15	8	16	99	-	-	-	5	2	83
Cycles with oocyte retrieval	-	37	19	11	8	13	96	-	-	-	5	2	59
Cycles with gamete transfer	-	37	19	11	8	13	94	-	-	-	5	2	57
Clinical pregnancies	-	12	4	4	2	3	33	-	-	-	3	-	10
Viable pregnancies	-	9	3	2	2	3	25	-	-	-	3	-	7
Clinical pregnancies per 100 oocyte retrieval cycles	-	32.4	21.1	36.4	25.0	23.1	34.4	-	-	-	60.0	-	16.9
Viable pregnancies per 100 oocyte retrieval cycles	-	24.3	15.8	18.2	25.0	23.1	26.0	-	-	-	60.0	-	11.9

Stage of treatment	IVF unit										Total
	AA	AB	AC	AD	AE	AF	AG	AH	AI		
Treatment cycles commenced	-	164	4	-	-	3	1	15	3		2,195
Cycles with oocyte retrieval	-	146	3	-	-	3	-	14	3		1,924
Cycles with gamete transfer	-	141	2	-	-	3	-	13	3		1,858
Clinical pregnancies	-	36	-	-	-	-	-	3	1		506
Viable pregnancies	-	27	-	-	-	-	-	1	1		384
Clinical pregnancies per 100 oocyte retrieval cycles	-	24.7	-	-	-	-	-	21.4	33.3		26.3
Viable pregnancies per 100 oocyte retrieval cycles	-	18.5	-	-	-	-	-	7.1	33.3		20.0

**Table 11: IVF pregnancies after embryo freezing but without ICSI, numbers and pregnancy rates in each IVF unit, 1997**

Stage of treatment	IVF unit												
	A	B	C	D	E	F	G	H	I	J	K	L	M
Cycles with embryo transfer	789	282	216	415	247	154	334	205	222	169	147	158	218
Clinical pregnancies	122	36	29	65	46	40	60	42	30	26	26	29	13
Viable pregnancies	95	31	23	50	28	33	59	37	25	17	20	19	8
Clinical pregnancies per 100 embryo transfers	15.5	12.8	13.4	15.7	18.6	26.0	18.0	20.5	13.5	15.4	17.7	18.4	6.0
Viable pregnancies per 100 embryo transfers	12.0	11.0	10.6	12.0	11.3	21.4	17.7	18.0	11.3	10.1	13.6	12.0	3.7

Stage of treatment	IVF unit												
	N	O	P	Q	R	S	T	U	V	W	X	Y	Z
Cycles with embryo transfer	119	85	133	78	100	85	17	45	21	8	30	38	19
Clinical pregnancies	17	13	10	12	31	11	1	8	2	1	6	5	3
Viable pregnancies	8	10	7	7	24	10	-	5	2	1	3	4	3
Clinical pregnancies per 100 embryo transfers	14.3	15.3	7.5	15.4	31.0	12.9	5.9	17.8	9.5	12.5	20.0	13.2	15.8
Viable pregnancies per 100 embryo transfers	6.7	11.8	5.3	9.0	24.0	11.8	-	11.1	9.5	12.5	10.0	10.5	15.8

Stage of treatment	IVF unit										Total
	AA	AB	AC	AD	AE	AF	AG	AH	AI		
Cycles with embryo transfer	52	-	12	20	34	38	17	5	8		4,520
Clinical pregnancies	9	-	-	5	7	5	4	-	1		715
Viable pregnancies	5	-	-	4	4	4	4	-	1		551
Clinical pregnancies per 100 embryo transfers	17.3	-	-	25.0	20.6	13.2	23.5	-	12.5		15.8
Viable pregnancies per 100 embryo transfers	9.6	-	-	20.0	11.8	10.5	23.5	-	12.5		12.2

**Table 12: IVF pregnancies after ICSI and embryo freezing, numbers and pregnancy rates in each IVF unit, 1997**

Stage of treatment	IVF unit												
	A	B	C	D	E	F	G	H	I	J	K	L	M
Cycles with embryo transfer	824	143	122	279	154	121	118	232	143	91	54	58	141
Clinical pregnancies	106	22	11	44	31	36	28	18	20	12	11	6	9
Viable pregnancies	85	19	6	39	24	31	24	14	18	6	10	2	6
Clinical pregnancies per 100 embryo transfers	12.9	15.4	9.0	15.8	20.1	29.8	23.7	7.8	14.0	13.2	20.4	10.3	6.4
Viable pregnancies per 100 embryo transfers	10.3	13.3	4.9	14.0	15.6	25.6	20.3	6.0	12.6	6.6	18.5	3.4	4.3

Stage of treatment	IVF unit												
	N	O	P	Q	R	S	T	U	V	W	X	Y	Z
Cycles with embryo transfer	39	149	89	111	56	19	18	73	10	15	9	14	6
Clinical pregnancies	6	13	4	24	19	2	2	13	-	1	2	1	1
Viable pregnancies	3	10	3	14	18	2	2	12	-	1	2	1	1
Clinical pregnancies per 100 embryo transfers	15.4	8.7	4.5	21.6	33.9	10.5	11.1	17.8	-	6.7	22.2	7.1	16.7
Viable pregnancies per 100 embryo transfers	7.7	6.7	3.4	12.6	32.1	10.5	11.1	16.4	-	6.7	22.2	7.1	16.7

Stage of treatment	IVF unit										Total
	AA	AB	AC	AD	AE	AF	AG	AH	AI		
Cycles with embryo transfer	40	-	-	20	16	11	27	1	-		3,203
Clinical pregnancies	7	-	-	3	4	-	1	-	-		457
Viable pregnancies	5	-	-	2	3	-	1	-	-		364
Clinical pregnancies per 100 embryo transfers	17.5	-	-	15.0	25.0	-	3.7	-	-		14.3
Viable pregnancies per 100 embryo transfers	12.5	-	-	10.0	18.8	-	3.7	-	-		11.4

**Table 13: Treatment related to embryo freezing in each IVF unit, 1997**

Stage of treatment	IVF unit												
	A	B	C	D	E	F	G	H	I	J	K	L	M
Patients having embryos frozen	957	600	417	551	301	445	350	296	231	197	243	122	167
Embryos that were frozen	4,540	2,067	1,966	2,729	1,081	2,115	2,115	1,594	1,457	1,013	1,158	964	989
Embryos thawed	3,340	1,490	1,295	2,003	1,019	1,087	1,296	1,316	987	646	722	671	839
Patients receiving thawed embryos	988	415	292	467	301	322	333	237	245	195	179	157	184
Embryos transferred after thawing	2,664	1,010	864	1,280	754	879	923	951	720	524	437	421	595
Frozen embryos in storage	8,597	4,620	2,019	4,748	na	na	2,997	3,460	2,690	1,631	1,560	1,413	1,133

Stage of treatment	IVF unit												
	N	O	P	Q	R	S	T	U	V	W	X	Y	Z
Patients having embryos frozen	111	187	151	153	142	121	91	39	44	40	41	74	49
Embryos that were frozen	725	668	848	862	1,099	608	372	731	234	309	271	322	207
Embryos thawed	475	609	663	598	616	343	188	475	171	121	156	301	130
Patients receiving thawed embryos	168	247	110	123	131	90	35	76	34	20	46	52	27
Embryos transferred after thawing	374	442	522	437	456	288	84	317	64	40	101	98	55
Frozen embryos in storage	933	1,170	1,116	1,116	1,818	917	546	529	266	306	366	483	322

Stage of treatment	IVF unit										Total
	AA	AB	AC	AD	AE	AF	AG	AH	AI		
Patients having embryos frozen	41	-	34	40	37	40	40	28	11		6,391
Embryos that were frozen	266	-	180	179	233	166	146	71	42		32,327
Embryos thawed	345	-	65	133	161	161	147	17	25		22,611
Patients receiving thawed embryos	64	-	12	40	32	49	34	6	8		5,719
Embryos transferred after thawing	199	-	27	84	105	116	99	14	15		15,959
Frozen embryos in storage	501	-	115	139	323	216	188	67	17		46,322 <sup>(a)</sup>

(a) Excludes E and F

**Table 14: Numbers and outcomes of IVF pregnancies by year of conception, 1979–1997**

Outcome of pregnancy	Year of conception							
	1979–95		1996		1997		1979–97	
	Number	Per cent	Number	Per cent	Number	Per cent	Number	Per cent
Spontaneous abortion	3,281	20.8	643	21.2	681	19.5	4,605	20.6
Termination of pregnancy	84	0.5	15	0.5	22	0.6	121	0.5
Ectopic pregnancy	712	4.5	93	3.1	91	2.6	896	4.0
Stillbirth	215	1.4	33	1.1	30	0.9	278	1.2
Live birth <sup>(a)</sup>	11,490	72.8	2,254	74.2	2,664	76.4	16,408	73.6
<b>All outcomes</b>	<b>15,782</b>	<b>100.0</b>	<b>3,038</b>	<b>100.0</b>	<b>3,488</b>	<b>100.0</b>	<b>22,308</b>	<b>100.0</b>

(a) Multiple pregnancies with both stillbirths and live births are included only in the live birth category

**Table 15: Place of parental residence, IVF pregnancies, 1979–1997**

Place of usual residence	Number			Per cent		
	1979–95	1996	1997	1979–95	1996	1997
	4,461	824	1,079	28.4	27.1	31.2
Victoria	4,439	928	919	28.2	30.6	26.6
Queensland	1,993	297	374	12.7	9.8	10.8
South Australia	1,892	304	303	12.0	10.0	8.8
Western Australia	1,332	270	336	8.5	8.9	9.7
Tasmania	295	62	101	1.9	2.0	2.9
Australian Capital Territory	138	35	40	0.9	1.2	1.2
Northern Territory	145	36	36	0.9	1.2	1.0
New Zealand	947	259	256	6.0	8.5	7.4
Other countries	84	21	15	0.5	0.7	0.4
Not stated	56	2	29			
<b>All regions</b>	<b>15,782</b>	<b>3,038</b>	<b>3,488</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

**Table 16: Maternal ages, IVF pregnancies, 1979–1997**

Age group (years)	Number			Per cent		
	1979–95	1996	1997	1979–95	1996	1997
Less than 20	4	-	-	0.0	-	-
20–24	340	55	71	2.2	1.8	2.0
25–29	3,619	548	608	23.0	18.0	17.4
30–34	6,797	1,197	1,375	43.1	39.4	39.4
35–39	4,198	964	1,137	26.6	31.7	32.6
40–44	754	233	265	4.8	7.7	7.6
45 and over	53	40	30	0.3	1.3	0.9
Not stated	17	1	2			
<b>All ages</b>	<b>15,782</b>	<b>3,038</b>	<b>3,488</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

**Table 17: Paternal ages, IVF pregnancies, 1979–1997**

Age group (years)	Number					
	1979–95	1996	1997	1979–95	1996	1997
Less than 20	1	-	-	0.0	-	-
20–24	141	9	20	0.9	0.3	0.6
25–29	2,036	339	316	13.7	11.2	9.1
30–34	5,506	927	1,077	37.0	30.7	31.1
35–39	4,441	935	1,168	29.9	30.9	33.8
40–44	1,867	522	543	12.6	17.3	15.7
45 and over	871	292	336	5.9	9.7	9.7
Not stated	919	14	28			
<b>All ages</b>	<b>15,782</b>	<b>3,038</b>	<b>3,488</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Table 18: Previous pregnancies for pregnant women, IVF pregnancies, 1979–1997

Number of previous pregnancies	Number				Per cent			
	1979–94	1995	1996	1997	1979–94	1995	1996	1997
One	6,107	1,348	1,612	1,816	47.9	51.9	53.1	52.2
Two	3,550	710	798	1,003	27.8	27.4	26.3	28.8
Three	1,651	293	357	395	12.9	11.3	11.8	11.4
Four or more	849	140	144	152	6.7	5.4	4.7	4.4
Not stated	604	104	123	112	4.7	4.0	4.1	3.2
	402	24	4	10				
<b>All women</b>	<b>13,163</b>	<b>2,619</b>	<b>3,038</b>	<b>3,488</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Table 19: Duration of infertility, IVF pregnancies, 1979–1997

Duration of infertility (years)	Number			Per cent		
	1979–95	1996	1997	1979–95	1996	1997
Less than 2	673	251	366	4.5	8.3	10.7
2–3	4,984	1,210	1,429	33.1	40.1	41.9
4–5	4,349	768	821	28.9	25.5	24.1
6–7	2,365	413	395	15.7	13.7	11.6
8–9	1,220	177	184	8.1	5.9	5.4
10 or more	1,452	195	215	9.7	6.5	6.3
Not stated	739	24	78			
<b>All pregnancies</b>	<b>15,782</b>	<b>3,038</b>	<b>3,488</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Table 20: Outcome of pregnancy by duration of infertility, IVF pregnancies, 1996–1997

Outcome of pregnancy	Duration of infertility (years)					
	Less than 4		4–7		8 or more	
	Number	Per cent	Number	Per cent	Number	Per cent
Spontaneous abortion	635	19.5	486	20.3	185	24.0
Termination of pregnancy	23	0.7	10	0.4	2	0.3
Ectopic pregnancy	85	2.6	73	3.0	24	3.1
Stillbirth	28	0.9	29	1.2	5	0.6
Live birth <sup>(a)</sup>	2,485	76.3	1,799	75.1	555	72.0
<b>All outcomes</b>	<b>3,256</b>	<b>100.0</b>	<b>2,397</b>	<b>100.0</b>	<b>771</b>	<b>100.0</b>

(a) Multiple pregnancies with both stillbirths and live births are included only in the live birth category  
 Note. The duration of infertility was not stated for 102 pregnancies

**Table 21: Causes of infertility, IVF pregnancies, 1979–1997**

Causes of infertility	Number			Per cent		
	1979–95	1996	1997	1979–95	1996	1997
Tubal	5,290	383	428	33.6	12.6	12.3
Male factor	3,031	1,040	1,158	19.2	34.3	33.3
Endometriosis	848	146	190	5.4	4.8	5.5
Other stated causes	848	225	320	5.4	7.4	9.2
Multiple causes	4,212	932	925	26.7	30.7	26.6
Unexplained infertility	1,535	306	456	9.7	10.1	13.1
Not stated	18	6	11			
<b>All causes</b>	<b>15,782</b>	<b>3,038</b>	<b>3,488</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

**Table 22: Outcome of pregnancy by causes of infertility, IVF pregnancies, 1996–1997**

Outcome of pregnancy	Causes of infertility					
	Tubal	Male	Endometriosis	Multiple	Unexplained	All causes <sup>(b)</sup>
	<b>Number</b>					
Spontaneous abortion	166	394	76	388	150	1,324
Termination of pregnancy	2	11	1	8	10	37
Ectopic pregnancy	45	42	6	50	24	184
Stillbirth	10	17	3	19	6	63
Live birth <sup>(a)</sup>	588	1,734	250	1,392	572	4,918
<b>All outcomes</b>	<b>811</b>	<b>2,198</b>	<b>336</b>	<b>1,857</b>	<b>762</b>	<b>6,526</b>
	<b>Per cent</b>					
Spontaneous abortion	20.5	17.9	22.6	20.9	19.7	20.3
Termination of pregnancy	0.2	0.5	0.3	0.4	1.3	0.6
Ectopic pregnancy	5.5	1.9	1.8	2.7	3.1	2.8
Stillbirth	1.2	0.8	0.9	1.0	0.8	1.0
Live birth <sup>(a)</sup>	72.5	78.9	74.4	75.0	75.1	75.4
<b>All outcomes</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Multiple pregnancies with both stillbirths and live births are included only in the live birth category

(b) Includes 562 pregnancies with 'other' or 'not stated' causes of infertility

**Table 23: Drugs used to stimulate ovulation, IVF pregnancies, 1979–1997**

Drugs	Number			Per cent		
	1979–95	1996	1997	1979–95	1996	1997
Natural cycles	164	133	145	1.1	4.4	4.2
Clomiphene and hMG or FSH	5,925	77	59	39.8	2.5	1.7
hMG or FSH	284	25	134	1.9	0.8	3.8
GnRH analogues and hMG or FSH	8,445	2,802	3,144	56.8	92.3	90.2
Other	59	-	3	0.4	-	0.1
Not stated	905	1	3			
<b>All drugs</b>	<b>15,782</b>	<b>3,038</b>	<b>3,488</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Table 24: IVF treatment cycle in which conception occurred, 1979–1997

Treatment cycle	Number			Per cent		
	1979–95	1996	1997	1979–95	1996	1997
1	6,522	1,330	1,698	43.5	43.9	48.9
2	3,653	782	839	24.4	25.8	24.2
3	2,072	418	432	13.8	13.8	12.4
4	1,137	222	218	7.6	7.3	6.3
5 or more	1,609	280	283	10.7	9.2	8.2
Not stated	789	6	18			
<b>All cycles</b>	<b>15,782</b>	<b>3,038</b>	<b>3,488</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Table 25: Number of oocytes collected by laparoscopy or ultrasound guidance, IVF pregnancies, 1979–1997

Number of oocytes collected	1979–95		1996		1997	
	Number	Per cent	Number	Per cent	Number	Per cent
1–2	860	5.9	95	3.2	96	2.8
3–4	2,192	15.0	223	7.6	231	6.8
5–6	2,699	18.5	365	12.4	363	10.7
7–8	2,824	19.4	420	14.3	467	13.7
9–10	1,522	10.4	360	12.3	474	13.9
11–12	1,243	8.5	349	11.9	427	12.5
13–14	932	6.4	314	10.7	352	10.3
15 or more	2,320	15.9	809	27.6	993	29.2
Not stated	1,190		103		85	
<b>All pregnancies</b>	<b>15,782</b>	<b>100.0</b>	<b>3,038</b>	<b>100.0</b>	<b>3,488</b>	<b>100.0</b>
Mean number of oocytes	9.9 (a)		11.6		12.0	

(a) Data include 1987–1995

Table 26: Number of embryos transferred, IVF pregnancies, 1979–1997

Number of embryos transferred	Number			Per cent		
	1979–95	1996	1997	1979–95	1996	1997
1	1,227	239	265	7.9	7.9	7.7
2	5,649	1,717	2,021	36.3	56.6	58.8
3	6,764	1,001	1,092	43.4	33.0	31.8
4	1,767	73	60	11.3	2.4	1.7
5 or more	167	4	1	1.1	0.1	0.0
Not stated	208	4	49			
<b>All pregnancies</b>	<b>15,782</b>	<b>3,038</b>	<b>3,488</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
Mean number of embryos	2.6	2.3	2.3			



**Table 27: Outcome of pregnancy by number of embryos transferred, IVF pregnancies, 1996–1997**

Outcome of pregnancy	Number of embryos transferred					All <sup>(b)</sup> pregnancies
	1	2	3	4	5+	
	<b>Number</b>					
Spontaneous abortion	115	732	427	36	1	1,324
Termination of pregnancy	-	27	6	3	-	37
Ectopic pregnancy	14	107	60	2	-	184
Stillbirth	5	36	19	2	-	63
Live birth <sup>(a)</sup>	370	2,836	1,581	90	4	4,918
<b>All outcomes</b>	<b>504</b>	<b>3,738</b>	<b>2,093</b>	<b>133</b>	<b>5</b>	<b>6,526</b>
	<b>Per cent</b>					
Spontaneous abortion	22.8	19.6	20.4	27.1	20.0	20.3
Termination of pregnancy	-	0.7	0.3	2.3	-	0.6
Ectopic pregnancy	2.8	2.9	2.9	1.5	-	2.8
Stillbirth	1.0	1.0	0.9	1.5	-	1.0
Live birth <sup>(a)</sup>	73.4	75.9	75.5	67.7	80.0	75.4
<b>All outcomes</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Multiple pregnancies with both stillbirths and live births are included only in the live birth category

(b) Includes 53 pregnancies in which the number of embryos transferred was not stated

**Table 28: Number of IVF pregnancies following donor oocytes, sperm or embryos, and frozen embryos or oocytes, 1979–1997**

Type of pregnancy	1979–94	1995	1996	1997	1979–97
Donor sperm	359	89	114	121	683
Donor embryos	1,122	135	131	128	1,516
Frozen embryos	29	15	22	12	78
Frozen oocytes	2,869	915	1,014	1,192	5,990
	4	2	3	1	10

**Table 29: Outcome of pregnancy after use of donor gametes, donor or frozen embryos, IVF pregnancies, 1979–1997**

Outcome of pregnancy	Donor sperm		Donor oocytes		Donor embryos		Frozen embryos	
	Number	Per cent	Number	Per cent	Number	Per cent	Number	Per cent
Spontaneous abortion	293	19.3	175	25.6	27	34.6	1,162	19.4
Termination of pregnancy	8	0.5	6	0.9	-	-	38	0.6
Ectopic pregnancy	45	3.0	12	1.8	2	2.6	165	2.8
Stillbirth	26	1.7	7	1.0	1	1.3	49	0.8
Live birth <sup>(a)</sup>	1,144	75.5	483	70.7	48	61.5	4,576	76.4
<b>All outcomes</b>	<b>1,516</b>	<b>100.0</b>	<b>683</b>	<b>100.0</b>	<b>78</b>	<b>100.0</b>	<b>5,990</b>	<b>100.0</b>

(a) Multiple pregnancies with both stillbirths and live births are included only in the live birth category

**Table 30: Number and outcome of pregnancies after microinsemination, 1990–1997**

Outcome of pregnancy	Year of conception							
	1990–95		1996		1997		1990–97	
	Number	Per cent	Number	Per cent	Number	Per cent	Number	Per cent
Spontaneous abortion	330	19.0	253	20.7	270	17.8	853	19.1
Termination of pregnancy	14	0.8	6	0.5	7	0.5	27	0.6
Ectopic pregnancy	31	1.8	27	2.2	27	1.8	85	1.9
Stillbirth	21	1.2	9	0.7	12	0.8	42	0.9
Live birth <sup>(a)</sup>	1,341	77.2	929	75.9	1,199	79.1	3,469	77.5
<b>All outcomes</b>	<b>1,737</b>	<b>100.0</b>	<b>1,224</b>	<b>100.0</b>	<b>1,515</b>	<b>100.0</b>	<b>4,476</b>	<b>100.0</b>

(a) Multiple pregnancies with both stillbirths and live births are included only in the live birth category

**Table 31: Drugs used in the luteal phase after embryo transfer, IVF pregnancies, 1979–1997**

Drugs	Number			Per cent		
	1979–95	1996	1997	1979–95	1996	1997
Proluton	666	579	700	4.3	19.1	20.1
Human chorionic gonadotrophin (hCG)	7,490	1,367	1,553	48.1	45.0	44.5
Human chorionic gonadotrophin / Proluton	726	232	213	4.7	7.6	6.1
Progestagen	2,221	285	393	14.2	9.4	11.3
Other drugs	87	10	30	0.6	0.3	0.9
None	4,396	565	599	28.2	18.6	17.2
Not stated	196	-	-			
<b>All pregnancies</b>	<b>15,782</b>	<b>3,038</b>	<b>3,488</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Table 32: Outcome of pregnancy by maternal age groups, IVF pregnancies, 1979–1997

Outcome of pregnancy	Maternal age (years)					All ages <sup>(b)</sup>
	Less than 25	25–29	30–34	35–39	40 and over	
	<b>Number</b>					
Spontaneous abortion	21	208	448	451	196	1,324
Termination of pregnancy	-	3	11	14	9	37
Ectopic pregnancy	1	34	81	51	17	184
Stillbirth	2	13	21	20	7	63
Live birth <sup>(a)</sup>	102	898	2,011	1,565	339	4,918
<b>All outcomes</b>	<b>126</b>	<b>1,156</b>	<b>2,572</b>	<b>2,101</b>	<b>568</b>	<b>6,526</b>
	<b>Per cent</b>					
Spontaneous abortion	16.7	18.0	17.4	21.5	34.5	20.3
Termination of pregnancy	-	0.3	0.4	0.7	1.6	0.6
Ectopic pregnancy	0.8	2.9	3.1	2.4	3.0	2.8
Stillbirth	1.6	1.1	0.8	1.0	1.2	1.0
Live birth <sup>(a)</sup>	81.0	77.7	78.2	74.5	59.7	75.4
<b>All outcomes</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Multiple pregnancies with both stillbirths and live births are included only in the live birth category

(b) Includes 3 pregnancies in which maternal age was not stated

Table 33: Spontaneous abortions, IVF pregnancies, 1979–1997

Outcome of pregnancy	1979–94	1995	1996	1997	1979–97
Spontaneous abortion	2,787	494	643	681	4,605
Stillbirth	182	33	33	30	278
Live birth	9,480	2,010	2,254	2,664	16,408
Total abortions and births	12,449	2,537	2,930	3,375	21,291
Spontaneous abortion rate (%)	22.4	19.5	21.9	20.2	21.6

Table 34: Incidence of spontaneous abortions in maternal age groups, IVF pregnancies, 1979–1997

Maternal age (years)	Number of IVF pregnancies <sup>(a)</sup>	Spontaneous abortions	
		Number	Per cent
	125	21	16.8
25–29	1,119	208	18.6
30–34	2,480	448	18.1
35–39	2,036	451	22.2
40–44	472	171	36.2
45 and over	70	25	35.7
Not stated	3	0	
<b>All ages</b>	<b>6,305</b>	<b>1,324</b>	<b>21.0</b>

(a) Spontaneous abortions and pregnancies of at least 20 weeks' gestation

**Table 35: Ectopic pregnancies after IVF, 1979–1997**

Outcome of pregnancy	1979–94	1995	1996	1997	1979–97
Ectopic pregnancies	648	64	93	91	896
Clinical pregnancies	13,163	2,619	3,038	3,488	22,308
% ectopic pregnancies	4.9	2.4	3.1	2.6	4.0
Total abortions and births	12,515	2,555	2,945	3,397	21,412
Ectopic pregnancy ratio <sup>(a)</sup>	1:19.3	1:39.9	1:31.7	1:37.3	1:23.9

(a) Ratio of ectopic pregnancies: total abortions and births

**Table 36: Heterotopic pregnancies after IVF, 1979–1997**

Type of heterotopic pregnancy	Year							1979–97
	1979–91	1992	1993	1994	1995	1996	1997	
	<b>Number</b>							
Abortion	47	5	4	2	5	8	4	75
Birth	20	3	5	4	6	5	2	45
<b>All heterotopic pregnancies</b>	<b>67</b>	<b>8</b>	<b>9</b>	<b>6</b>	<b>11</b>	<b>13</b>	<b>6</b>	<b>120</b>
All pregnancies	7,683	1,513	1,708	2,259	2,619	3,038	3,488	22,308
	<b>Per cent</b>							
Abortion	0.6	0.3	0.2	0.1	0.2	0.3	0.1	0.3
Birth	0.3	0.2	0.3	0.2	0.2	0.2	0.1	0.2
<b>All heterotopic pregnancies</b>	<b>0.9</b>	<b>0.5</b>	<b>0.5</b>	<b>0.3</b>	<b>0.4</b>	<b>0.4</b>	<b>0.2</b>	<b>0.5</b>

**Table 37: Reported complications of pregnancy, IVF pregnancies, 1990–1997**

Pregnancy complications	Number			Per cent		
	1990–95	1996	1997	1990–95	1996	1997
None	5,829	2,254	2,520	67.4	74.3	74.8
Threatened abortion	471	121	116	5.4	4.0	3.4
Antepartum haemorrhage	163	41	60	1.9	1.4	1.8
Pregnancy-induced hypertension	535	195	208	6.2	6.4	6.2
Placenta praevia	104	32	38	1.2	1.1	1.1
Other complications	1,542	392	425	17.8	12.9	12.6
Not stated	2,007	3	121			
<b>All pregnancies</b>	<b>10,651</b>	<b>3,038</b>	<b>3,488</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

**Table 38: Duration of singleton and multiple IVF pregnancies of at least 20 weeks' gestation, 1996–1997**

Gestational age (weeks)	Singleton		Twin		Triplet		All pregnancies <sup>(a)</sup>	
	Number	Per cent	Number	Per cent	Number	Per cent	Number	Per cent
20–23	41	1.0	20	2.2	2	2.8	63	1.3
24–27	29	0.7	32	3.6	4	5.6	65	1.3
28–31	59	1.5	82	9.1	23	32.4	167	3.4
32–36	405	10.3	431	47.9	39	54.9	876	17.8
37–41	3,379	85.6	330	36.7	3	4.2	3,712	75.4
42 or more	36	0.9	4	0.4	-	-	40	0.8
20–36	534	13.5	565	62.8	68	95.8	1,171	23.8
Not stated	52		6		-		58	
<b>All gestational ages</b>	<b>4,001</b>	<b>100.0</b>	<b>905</b>	<b>100.0</b>	<b>71</b>	<b>100.0</b>	<b>4,981</b>	<b>100.0</b>

(a) Includes 4 quadruplet pregnancies

**Table 39: Duration of singleton and multiple IVF pregnancies of at least 20 weeks' gestation after microinsemination, 1990–1997**

Gestational age (weeks)	Singleton		Twin		Triplet		All pregnancies <sup>(a)</sup>	
	Number	Per cent	Number	Per cent	Number	Per cent	Number	Per cent
20–23	16	0.6	15	2.4	4	8.5	35	1.0
24–27	17	0.6	29	4.7	1	2.1	47	1.4
28–31	35	1.2	39	6.3	17	36.2	95	2.7
32–36	268	9.6	280	45.5	24	51.1	573	16.5
37–41	2,442	87.0	247	40.2	1	2.1	2,690	77.5
42 or more	28	1.0	5	0.8	-	-	33	1.0
20–36	336	12.0	363	59.0	46	97.9	750	21.6
Not stated	36		2		-		38	
<b>All gestational ages</b>	<b>2,842</b>	<b>100.0</b>	<b>617</b>	<b>100.0</b>	<b>47</b>	<b>100.0</b>	<b>3,511</b>	<b>100.0</b>

(a) Include 5 quadruplet pregnancies

**Table 40: Duration of pregnancy of singleton IVF births after use of donor sperm, donor oocytes or frozen embryos, 1996–1997**

Gestational age (weeks)	Donor sperm		Donor oocytes		Frozen embryos	
	Number	Per cent	Number	Per cent	Number	Per cent
20–27	4	2.9	4	3.1	22	1.5
28–31	2	1.5	1	0.8	19	1.3
32–36	13	9.6	21	16.0	131	9.1
37–41	117	86.0	105	80.2	1,263	87.3
42 or more	-	-	-	-	11	0.8
Not stated	3		-		17	
<b>All gestational ages</b>	<b>139</b>	<b>100.0</b>	<b>131</b>	<b>100.0</b>	<b>1,463</b>	<b>100.0</b>

**Table 41: Maternal age and duration of singleton IVF pregnancies of at least 20 weeks' gestation, 1996–1997**

Gestational age (weeks)	Maternal age (years)					All ages <sup>(a)</sup>
	Less than 25	25–29	30–34	35–39	40 and over	
	<b>Number</b>					
20–27	3	11	24	24	8	70
28–31	1	10	16	26	6	59
32–36	11	62	150	148	34	405
37 or more	65	615	1,364	1,109	261	3,415
Not stated	2	14	18	14	3	52
<b>All gestational ages</b>	<b>82</b>	<b>712</b>	<b>1,572</b>	<b>1,321</b>	<b>312</b>	<b>4,001</b>
20–36	15	83	190	198	48	534
	<b>Per cent</b>					
20–27	3.8	1.6	1.5	1.8	2.6	1.8
28–31	1.3	1.4	1.0	2.0	1.9	1.5
32–36	13.8	8.9	9.7	11.3	11.0	10.3
37 or more	81.3	88.1	87.8	84.9	84.5	86.5
<b>All gestational ages</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
20–36	18.8	11.9	12.2	15.1	15.5	13.5

(a) Includes 2 pregnancies in which maternal age was not stated

**Table 42: Causes of infertility and duration of singleton IVF pregnancies of at least 20 weeks' gestation, 1996-1997**

Gestational age (weeks)	Causes of infertility					
	Tubal	Male	Endometriosis	Multiple	Unexplained	All causes <sup>(a)</sup>
	Number					
20-27	12	17	4	24	6	70
28-31	7	23	2	10	9	59
32-36	52	130	14	126	44	405
37 or more	405	1,201	170	986	391	3,415
Not stated	6	25	3	7	5	52
<b>All gestational ages</b>	<b>482</b>	<b>1,396</b>	<b>193</b>	<b>1,153</b>	<b>455</b>	<b>4,001</b>
20-36	71	170	20	160	59	534
	Per cent					
20-27	2.5	1.2	2.1	2.1	1.3	1.8
28-31	1.5	1.7	1.1	0.9	2.0	1.5
32-36	10.9	9.5	7.4	11.0	9.8	10.3
37-or more	85.1	87.6	89.5	86.0	86.9	86.5
<b>All gestational ages</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
20-36	14.9	12.4	10.5	14.0	13.1	13.5

(a) Includes 322 pregnancies with 'other' or 'not stated' causes of infertility

**Table 43: Plurality of IVF pregnancies of at least 20 weeks' gestation, 1979-1997**

Plurality	Number			Per cent		
	1979-95	1996	1997	1979-95	1996	1997
Singletons	9,424	1,852	2,149	80.5	81.0	79.8
Twins	2,011	399	506	17.2	17.4	18.8
Triplets	259	32	39	2.2	1.4	1.4
Quadruplets	10	4	-	0.1	0.2	-
Quintuplets	1	-	-	0.0	-	-
<b>All pregnancies</b>	<b>11,705</b>	<b>2,287</b>	<b>2,694</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

**Table 44: Plurality of IVF pregnancies of at least 20 weeks' gestation and number of embryos transferred, 1996–1997**

Number of embryos transferred	Plurality							
	Singleton		Twin		Triplet		All pregnancies <sup>(a)</sup>	
	Number	Per cent	Number	Per cent	Number	Per cent	Number	Per cent
1	371	98.9	3	0.8	1	0.3	375	100.0
2	2,329	81.1	538	18.7	5	0.2	2,872	100.0
3	1,193	74.6	341	21.3	62	3.9	1,600	100.0
4	72	78.3	17	18.5	3	3.3	92	100.0
5 or more	3	75.0	1	25.0	-	-	4	100.0
Not stated	33		5		-		38	
<b>All pregnancies</b>	<b>4,001</b>	<b>80.3</b>	<b>905</b>	<b>18.2</b>	<b>71</b>	<b>1.4</b>	<b>4,981</b>	<b>100.0</b>

(a) Includes 4 quadruplet pregnancies

**Table 45: Multiple pregnancies, IVF and GIFT pregnancies, State and Territory, 1996–1997**

Plurality	IVF Unit							Australia	New Zealand
	NSW	Vic	Qld	WA	SA/NT	ACT/Tas			
	<b>Number</b>								
Singletons	1,432	1,281	616	395	459	173	4,356	325	
Twins	284	291	162	100	116	41	994	88	
Triplets	28	22	24	9	2	1	86	9	
Quadruplets	1	-	1	3	-	-	5	-	
All pregnancies	1,745	1,594	803	507	577	215	5,441	422	
	<b>Per cent of viable pregnancies</b>								
Twins	16.3	18.3	20.2	19.7	20.1	19.1	18.3	20.9	
Triplets / Quadruplets	1.7	1.4	3.1	2.4	0.3	0.5	1.7	2.1	



**Table 46: Multiple pregnancies in each IVF unit, IVF and GIFT pregnancies, 1996–1997**

Plurality	IVF unit												
	A	B	C	D	E	F	G	H	I	J	K	L	M
	Number												
Singletons	671	534	427	416	243	315	228	206	149	126	158	96	81
Twins	120	155	104	105	49	90	47	33	25	16	54	15	16
Triplets	8	18	19	2	4	7	1	1	1	2	3	2	-
Quadruplets	-	-	-	-	-	-	-	-	-	1	-	-	-
All pregnancies	799	707	550	523	296	412	276	240	175	145	215	113	97
	Per cent of viable pregnancies												
Twins	15.0	21.9	18.9	20.1	16.6	21.8	17.0	13.8	14.3	11.0	25.1	13.3	16.5
Triplets / Quadruplets	1.0	2.5	3.5	0.4	1.4	1.7	0.4	0.4	0.6	2.1	1.4	1.8	-

Plurality	IVF unit												
	N	O	P	Q	R	S	T	U	V	W	X	Y	Z
	Number												
Singletons	102	82	67	78	116	75	55	43	20	51	37	43	32
Twins	28	5	13	26	41	20	16	9	2	12	13	11	10
Triplets	8	1	2	1	5	-	3	-	2	3	2	0	1
Quadruplets	-	-	-	-	3	-	1	-	-	0	0	0	0
All pregnancies	138	88	82	105	165	95	75	52	24	66	52	54	43
	Per cent of viable pregnancies												
Twins	20.3	5.7	15.9	24.8	24.8	21.1	21.3	17.3	8.3	18.2	25.0	20.4	23.3
Triplets / Quadruplets	5.8	1.1	2.4	1.0	4.8	-	5.3	-	8.3	4.5	3.8	-	2.3

Plurality	IVF unit										Total
	AA	AB	AC	AD	AE	AF	AG	AH	AI		
	Number										
Singletons	63	49	12	25	27	17	22	12	3		4,681
Twins	16	9	2	3	7	5	-	4	1		1,082
Triplets	2	3	-	-	-	1	-	-	-		102
Quadruplets	-	-	-	-	-	-	-	-	-		5
All pregnancies	81	61	14	28	34	23	22	16	4		5,870
	Per cent of viable pregnancies										
Twins	19.8	14.8	14.3	10.7	20.6	21.7	-	25.0	25.0		18.4
Triplets / Quadruplets	2.5	4.9	-	-	-	4.3	-	-	-		1.8

**Table 47: Method of delivery for singleton and multiple IVF pregnancies of at least 20 weeks' gestation, 1996–1997**

	Method of delivery				
	Vaginal		Caesarean section		All methods <sup>(a)</sup>
	Number	Per cent	Number	Per cent	Number
Singleton	2,548	64.2	1,421	35.8	4,001
Twin	349	39.0	547	61.0	905
Triplet	9	12.7	62	87.3	71
Quadruplet	-	-	4	100.0	4
<b>All pregnancies</b>	<b>2,906</b>	<b>58.8</b>	<b>2,034</b>	<b>41.2</b>	<b>4,981</b>

(a) Includes 41 pregnancies in which the method of delivery was not stated

**Table 48: Sex of infants in singleton and multiple IVF births of at least 20 weeks' gestation, 1979–1997**

Plurality	Male			Female			Sex ratio (M:F)		
	1979–95	1996	1997	1979–95	1996	1997	1979–95	1996	1997
Singletons	4,838	960	1,112	4,541	887	1,030	106.5	108.2	108.0
Twins	2,065	404	529	1,937	393	481	106.6	102.8	110.0
Triplets	386	45	62	384	51	55	100.5	88.2	112.7
Quadruplets	23	7	-	17	9	-	135.3	77.8	-
Quintuplets	1	-	-	4	-	-	25.0	-	-
<b>All births</b>	<b>7,313</b>	<b>1,416</b>	<b>1,703</b>	<b>6,883</b>	<b>1,340</b>	<b>1,566</b>	<b>106.2</b>	<b>105.7</b>	<b>108.7</b>

Note. Infant's sex was not stated or indeterminate for 87 births

**Table 49: Birthweight of IVF live births and stillbirths, 1996–1997**

Birthweight (g)	Live births		Stillbirths		All births <sup>(a)</sup>	
	Number	Per cent	Number	Per cent	Number	Per cent
Less than 500	14	0.2	44	57.1	59	1.0
500–999	98	1.7	16	20.8	117	1.9
1000–1499	234	4.1	6	7.8	243	4.0
1500–1999	386	6.8	3	3.9	407	6.8
2000–2499	771	13.6	2	2.6	807	13.4
2500–2999	1,174	20.7	2	2.6	1,233	20.5
3000–3499	1,604	28.2	1	1.3	1,672	27.8
3500–3999	1,049	18.5	2	2.6	1,100	18.3
4000 and over	348	6.1	1	1.3	368	6.1
Not stated	16		12		34	
<b>All birthweights</b>	<b>5,694</b>	<b>100.0</b>	<b>89</b>	<b>100.0</b>	<b>6,040</b>	<b>100.0</b>
Mean birthweight (g)	2,924		816		2,900	

(a) Includes 257 infants with unstated outcome

**Table 50: Birthweight of infants in singleton and multiple IVF births of at least 20 weeks' gestation, 1996–1997**

Birthweight (g)	Singleton		Twin		Triplet		All births <sup>(a)</sup>	
	Number	Per cent	Number	Per cent	Number	Per cent	Number	Per cent
Less than 500	29	0.7	26	1.4	2	0.9	59	1.0
500–999	35	0.9	65	3.6	17	8.0	117	1.9
1000–1499	43	1.1	131	7.3	60	28.2	243	4.0
1500–1999	82	2.1	246	13.7	75	35.2	407	6.8
2000–2499	231	5.8	530	29.4	45	21.1	807	13.4
2500–2999	657	16.5	563	31.2	13	6.1	1,233	20.5
3000–3499	1,451	36.5	220	12.2	1	0.5	1,672	27.8
3500–3999	1,079	27.1	21	1.2	-	-	1,100	18.3
4000 and over	368	9.3	-	-	-	-	368	6.1
Less than 2500	420	10.6	998	55.4	199	93.4	1,633	27.2
Not stated	26		8		-		34	
<b>All birthweights</b>	<b>4,001</b>	<b>100.0</b>	<b>1,810</b>	<b>100.0</b>	<b>213</b>	<b>100.0</b>	<b>6,040</b>	<b>100.0</b>
Mean birthweight (g)	3,236		2,312		1,700		2,900	

(a) Includes 16 quadruplet births

**Table 51: Birthweight of infants in singleton and multiple births of at least 20 weeks' gestation after microinsemination, 1990–1997**

Birthweight (g)	Singleton		Twin		Triplet		All births <sup>(a)</sup>	
	Number	Per cent	Number	Per cent	Number	Per cent	Number	Per cent
Less than 500	11	0.4	19	1.6	8	5.7	40	1.0
500–999	24	0.8	66	5.4	9	6.4	99	2.4
1000–1499	26	0.9	75	6.1	45	31.9	159	3.8
1500–1999	58	2.1	146	11.9	42	29.8	250	5.9
2000–2499	160	5.7	363	29.7	28	19.9	552	13.1
2500–2999	486	17.2	375	30.7	7	5.0	868	20.6
3000–3499	1,061	37.5	165	13.5	2	1.4	1,228	29.2
3500–3999	737	26.1	13	1.1	-	-	750	17.8
4000 and over	264	9.3	-	-	-	-	264	6.3
Less than 2500	279	9.9	669	54.7	132	93.6	1,100	26.1
Not stated	15		12		-		27	
<b>All birthweights</b>	<b>2,842</b>	<b>100.0</b>	<b>1,234</b>	<b>100.0</b>	<b>141</b>	<b>100.0</b>	<b>4,237</b>	<b>100.0</b>
Mean birthweight (g)	3,248		2,307		1,623		2,911	

(a) Includes 20 quadruplet births

**Table 52: Birthweight of infants in singleton IVF births after use of donor sperm, donor oocytes or frozen embryos, 1979–1997**

Birthweight (g)	Donor sperm		Donor oocytes		Frozen embryos	
	Number	Per cent	Number	Per cent	Number	Per cent
Less than 500	8	0.9	4	1.0	16	0.4
500–999	10	1.1	5	1.2	40	1.0
1000–1499	12	1.3	8	1.9	34	0.9
1500–1999	15	1.6	9	2.2	57	1.4
2000–2499	51	5.6	29	7.0	157	3.9
2500 or more	814	89.5	361	86.8	3,675	92.4
Not stated	10		4		35	
Less than 2500	96	10.5	55	13.2	304	7.6
<b>All birthweights</b>	<b>920</b>	<b>100.0</b>	<b>420</b>	<b>100.0</b>	<b>4,014</b>	<b>100.0</b>
Mean birthweight (g)	3,236		3,209		3,342	

**Table 53: Outcome of infants in singleton and multiple IVF births of at least 20 weeks' gestation, 1996–1997**

Outcome	Singleton	Twin	Triplet	Quadruplet	All births <sup>(a)</sup>
Live births	3,948	1,778	211	14	5,951
Stillbirths	53	32	2	2	89
<b>Total births</b>	<b>4,001</b>	<b>1,810</b>	<b>213</b>	<b>16</b>	<b>6,040</b>
Neonatal deaths	18	30	9	-	57
Perinatal deaths	71	62	11	2	146
Stillbirth rate per 1,000 total births	13.2	17.7	9.4	125.0	14.7
Neonatal death rate per 1,000 live births	4.6	16.9	42.7	-	9.6
Perinatal mortality rate per 1,000 total births	17.7	34.3	51.6	125.0	24.2

(a) Live births include 257 births for which birth status was not recorded

**Table 54: Major congenital malformations in singleton and multiple IVF births of at least 20 weeks' gestation, 1979–1997**

Outcome	Singleton	Multiple	All births <sup>(a)</sup>
Total births	13,487	6,889	20,376
Congenital malformations			
— number	409	120	529
— rate (per cent)	3.0	1.7	2.6

(a) Includes 68 abortions for fetal abnormality at gestational ages of at least 16 weeks

**Table 55: Numbers and outcomes of GIFT pregnancies by year of conception, 1985–1997**

Outcome of pregnancy	Year of conception							
	1985–95		1996		1997		1985–97	
	Number	Per cent	Number	Per cent	Number	Per cent	Number	Per cent
Spontaneous abortion	1,580	20.7	127	19.4	115	22.8	1,822	20.7
Termination of pregnancy	45	0.6	1	0.2	7	1.4	53	0.6
Ectopic pregnancy	256	3.4	11	1.7	16	3.2	283	3.2
Stillbirth	84	1.1	3	0.5	6	1.2	93	1.1
Live birth <sup>(a)</sup>	5,669	74.3	512	78.3	361	71.5	6,542	74.4
<b>All outcomes</b>	<b>7,634</b>	<b>100.0</b>	<b>654</b>	<b>100.0</b>	<b>505</b>	<b>100.0</b>	<b>8,793</b>	<b>100.0</b>

(a) Multiple pregnancies with both stillbirths and live births are included only in the live birth category

**Table 56: Place of parental residence, GIFT pregnancies, 1985–1997**

Place of usual residence	Number			Per cent		
	1985–95	1996	1997	1985–95	1996	1997
New South Wales	2,469	254	156	32.4	38.8	31.0
Victoria	1,577	101	79	20.7	15.4	15.7
Queensland	2,176	238	221	28.5	36.4	43.8
South Australia	484	25	20	6.3	3.8	4.0
Western Australia	674	23	13	8.8	3.5	2.6
Tasmania	30	2	-	0.4	0.3	-
Australian Capital Territory	51	6	3	0.7	0.9	0.6
Northern Territory	18	2	1	0.2	0.3	0.2
New Zealand	115	2	7	1.5	0.3	1.4
Other countries	29	1	4	0.4	0.2	0.8
Not stated	11	-	1			
<b>All regions</b>	<b>7,634</b>	<b>654</b>	<b>505</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

**Table 57: Maternal ages, GIFT pregnancies, 1985–1997**

Age group (years)	Number			Per cent		
	1985–95	1996	1997	1985–95	1996	1997
Less than 20	-	-	-	-	-	-
20–24	199	7	4	2.6	1.1	0.8
25–29	1,833	128	80	24.0	19.6	15.8
30–34	3,306	272	213	43.3	41.6	42.2
35–39	1,896	193	157	24.8	29.5	31.1
40–44	379	53	51	5.0	8.1	10.1
45 and over	19	1	-	0.2	0.2	-
Not stated	2	-	-			
<b>All ages</b>	<b>7,634</b>	<b>654</b>	<b>505</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Table 58: Paternal ages, GIFT pregnancies, 1985–1997

Age group (years)	Number			Per cent		
	1985–95	1996	1997	1985–95	1996	1997
Less than 20	2	-	-	0.0	-	-
20–24	38	2	2	0.5	0.3	0.4
25–29	1,033	82	43	14.1	12.6	8.6
30–34	2,801	230	179	38.3	35.3	35.9
35–39	2,085	207	175	28.5	31.8	35.1
40–44	902	86	66	12.3	13.2	13.3
45 and over	448	44	33	6.1	6.8	6.6
Not stated	325	3	7			
<b>All ages</b>	<b>7,634</b>	<b>654</b>	<b>505</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Table 59: Previous pregnancies for pregnant women, GIFT pregnancies, 1985–1997

Number of previous pregnancies	Number				Per cent			
	1985–94	1995	1996	1997	1985–94	1995	1996	1997
None	3,707	320	348	275	54.3	49.5	53.2	54.9
One	2,078	203	180	144	30.5	31.4	27.5	28.7
Two	672	76	75	63	9.8	11.7	11.5	12.6
Three	237	29	31	13	3.5	4.5	4.7	2.6
Four or more	129	19	20	6	1.9	2.9	3.1	1.2
Not stated	157	7	-	4				
<b>All women</b>	<b>6,980</b>	<b>654</b>	<b>654</b>	<b>505</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Table 60: Duration of infertility, GIFT pregnancies, 1985–1997

Duration of infertility (years)	Number			Per cent		
	1985–95	1996	1997	1985–95	1996	1997
Less than 2	508	65	50	6.9	10.0	10.2
2–3	3,027	289	226	41.3	44.4	46.1
4–5	1,928	184	142	26.3	28.3	29.0
6–7	932	55	41	12.7	8.4	8.4
8–9	484	31	15	6.6	4.8	3.1
10 or more	449	27	16	6.1	4.1	3.3
Not stated	306	3	15			
<b>All pregnancies</b>	<b>7,634</b>	<b>654</b>	<b>505</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

**Table 61: Outcome of pregnancy by duration of infertility, GIFT pregnancies, 1996–1997**

Outcome of pregnancy	Duration of infertility (years)					
	Less than 4		4–7		8 or more	
	Number	Per cent	Number	Per cent	Number	Per cent
Spontaneous abortion	132	21.0	78	18.5	25	28.1
Termination of pregnancy	7	1.1	1	0.2	-	-
Ectopic pregnancy	11	1.7	13	3.1	2	2.2
Stillbirth	2	0.3	7	1.7	-	-
Live birth <sup>(a)</sup>	478	75.9	323	76.5	62	69.7
<b>All outcomes</b>	<b>630</b>	<b>100.0</b>	<b>422</b>	<b>100.0</b>	<b>89</b>	<b>100.0</b>

(a) Multiple pregnancies with both stillbirths and live births are included only in the live birth category

Note. The duration of infertility was not stated for 18 pregnancies

**Table 62: Causes of infertility, GIFT pregnancies, 1985–1997**

Causes of infertility	Number			Per cent		
	1985–95	1996	1997	1985–95	1996	1997
Tubal	366	16	13	4.8	2.4	2.6
Male factor	1,559	111	79	20.5	17.0	15.6
Endometriosis	1,094	82	94	14.4	12.5	18.6
Other stated causes	630	97	62	8.3	14.8	12.3
Multiple causes	1,753	137	104	23.0	20.9	20.6
Unexplained infertility	2,221	211	153	29.1	32.3	30.3
Not stated	11	-	-			
<b>All causes</b>	<b>7,634</b>	<b>654</b>	<b>505</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Table 63: Outcome of pregnancy by causes of infertility, GIFT pregnancies, 1996–1997

Outcome of pregnancy	Causes of infertility					
	Tubal	Male	Endometriosis	Multiple	Unexplained	All causes <sup>(b)</sup>
	<b>Number</b>					
Spontaneous abortion	6	34	33	44	82	242
Termination of pregnancy	-	1	2	-	4	8
Ectopic pregnancy	2	1	1	6	9	27
Stillbirth	-	3	-	2	3	9
Live birth <sup>(a)</sup>	21	151	140	189	266	873
<b>All outcomes</b>	<b>29</b>	<b>190</b>	<b>176</b>	<b>241</b>	<b>364</b>	<b>1,159</b>
	<b>Per cent</b>					
Spontaneous abortion	20.7	17.9	18.8	18.3	22.5	20.9
Termination of pregnancy	-	0.5	1.1	-	1.1	0.7
Ectopic pregnancy	6.9	0.5	0.6	2.5	2.5	2.3
Stillbirth	-	1.6	-	0.8	0.8	0.8
Live birth <sup>(a)</sup>	72.4	79.5	79.5	78.4	73.1	75.3
<b>All outcomes</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Multiple pregnancies with both stillbirths and live births are included only in the live birth category

(b) Includes 159 pregnancies with 'other' or 'not stated' causes of infertility

Table 64: Drugs used to stimulate ovulation, GIFT pregnancies, 1985–1997

Drugs	Number			Per cent		
	1985–95	1996	1997	1985–95	1996	1997
Natural cycles	13	4	7	0.2	0.6	1.4
Clomiphene and hMG or FSH	2,836	9	10	37.5	1.4	2.0
hMG or FSH	155	5	10	2.0	0.8	2.0
GnRH analogues and hMG or FSH	4,548	636	477	60.1	97.2	94.5
Other	11	-	1	0.1	-	0.2
Not stated	71	-	-			
<b>All drugs</b>	<b>7,634</b>	<b>654</b>	<b>505</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Table 65: GIFT treatment cycle in which conception occurred, 1985–1997

Treatment cycle	Number			Per cent		
	1985–95	1996	1997	1985–95	1996	1997
1	3,557	350	253	47.2	53.5	50.4
2	1,901	151	122	25.2	23.1	24.3
3	975	76	73	12.9	11.6	14.5
4	506	41	30	6.7	6.3	6.0
5 or more	603	36	24	8.0	5.5	4.8
Not stated	92	-	3			
<b>All cycles</b>	<b>7,634</b>	<b>654</b>	<b>505</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>



**Table 66: Number of oocytes collected by laparoscopy or ultrasound guidance, GIFT pregnancies, 1985–1997**

Number of oocytes collected	1979–95		1996		1997	
	Number	Per cent	Number	Per cent	Number	Per cent
1–2	376	5.0	31	4.7	38	7.6
3–4	1,531	20.4	110	16.8	64	12.8
5–6	1,663	22.1	139	21.3	96	19.2
7–8	1,628	21.6	113	17.3	99	19.8
9–10	838	11.1	78	11.9	65	13.0
11–12	563	7.5	78	11.9	44	8.8
13–14	346	4.6	41	6.3	33	6.6
15 or more	578	7.7	63	9.6	61	12.2
Not stated	111		1		5	
<b>All pregnancies</b>	<b>7,634</b>	<b>100.0</b>	<b>654</b>	<b>100.0</b>	<b>505</b>	<b>100.0</b>
Mean number of oocytes	7.7 <sup>(a)</sup>		8.4		8.7	

(a) Data include 1987–1995

**Table 67: Number of oocytes transferred, GIFT pregnancies, 1985–1997**

Number of oocytes transferred	Number			Per cent		
	1985–95	1996	1997	1985–95	1996	1997
1	110	12	10	1.5	1.8	2.0
2	1,474	277	226	19.5	42.5	45.1
3	4,261	319	232	56.3	49.0	46.3
4	1,558	40	31	20.6	6.1	6.2
5 or more	160	3	2	2.1	0.5	0.4
Not stated	71	3	4			
<b>All pregnancies</b>	<b>7,634</b>	<b>654</b>	<b>505</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
Mean number of oocytes	3.0	2.6	2.6			

Table 68: Outcome of pregnancy by number of oocytes transferred, GIFT pregnancies, 1996–1997

Outcome of pregnancy	Number of oocytes transferred					All <sup>(b)</sup> pregnancies
	1	2	3	4	5+	
	<b>Number</b>					
Spontaneous abortion	9	102	104	27	-	242
Termination of pregnancy	1	3	3	-	-	8
Ectopic pregnancy	-	11	14	1	-	27
Stillbirth	-	3	5	1	-	9
Live birth <sup>(a)</sup>	12	384	425	42	5	873
<b>All outcomes</b>	<b>22</b>	<b>503</b>	<b>551</b>	<b>71</b>	<b>5</b>	<b>1,159</b>
	<b>Per cent</b>					
Spontaneous abortion	40.9	20.3	18.9	38.0	-	20.9
Termination of pregnancy	4.5	0.6	0.5	-	-	0.7
Ectopic pregnancy	-	2.2	2.5	1.4	-	2.3
Stillbirth	-	0.6	0.9	1.4	-	0.8
Live birth <sup>(a)</sup>	54.5	76.3	77.1	59.2	100.0	75.3
<b>All outcomes</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Multiple pregnancies with both stillbirths and live births are included only in the live birth category

(b) Includes 7 pregnancies in which the number of oocytes transferred was not stated

Table 69: Drugs used in the luteal phase after embryo transfer, GIFT pregnancies, 1985–1997

Drugs	Number			Per cent		
	1985–95	1996	1997	1985–95	1996	1997
Proluton	131	85	91	1.7	13.0	18.0
Human chorionic gonadotrophin (hCG)	4,943	488	349	65.3	74.6	69.1
Human chorionic gonadotrophin / Proluton	307	24	16	4.1	3.7	3.2
Progestagen	882	28	34	11.6	4.3	6.7
Other drugs	6	22	1	0.1	3.4	0.2
None	1,303	7	14	17.2	1.1	2.8
Not stated	62	-	-			
<b>All pregnancies</b>	<b>7,634</b>	<b>654</b>	<b>505</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

**Table 70: Outcome of pregnancy by maternal age groups, GIFT pregnancies, 1985–1997**

Outcome of pregnancy	Maternal age (years)					All ages <sup>(b)</sup>
	Less than 25	25–29	30–34	35–39	40 and over	
	<b>Number</b>					
Spontaneous abortion	34	305	718	550	215	1,822
Termination of pregnancy	1	8	17	13	14	53
Ectopic pregnancy	8	68	125	70	12	283
Stillbirth	2	17	37	33	4	93
Live birth <sup>(a)</sup>	165	1,643	2,894	1,580	258	6,542
<b>All outcomes</b>	<b>210</b>	<b>2,041</b>	<b>3,791</b>	<b>2,246</b>	<b>503</b>	<b>8,793</b>
	<b>Per cent</b>					
Spontaneous abortion	16.2	14.9	18.9	24.5	42.7	20.7
Termination of pregnancy	0.5	0.4	0.4	0.6	2.8	0.6
Ectopic pregnancy	3.8	3.3	3.3	3.1	2.4	3.2
Stillbirth	1.0	0.8	1.0	1.5	0.8	1.1
Live birth <sup>(a)</sup>	78.6	80.5	76.3	70.3	51.3	74.4
<b>All outcomes</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

(a) Multiple pregnancies with both stillbirths and live births are included only in the live birth category

(b) Includes 2 pregnancies in which maternal age was not stated

**Table 71: Spontaneous abortions, GIFT pregnancies, 1985–1997**

Outcome of pregnancy	1985–94	1995	1996	1997	1985–97
Spontaneous abortion	1,463	117	127	115	1,822
Stillbirth	76	8	3	6	93
Live birth	5,162	507	512	361	6,542
Total abortions and births	6,701	632	642	482	8,457
Spontaneous abortion rate (%)	21.8	18.5	19.8	23.9	21.5

**Table 72: Incidence of spontaneous abortions in maternal age groups, GIFT pregnancies, 1985–1997**

Maternal age (years)	Number of GIFT pregnancies <sup>(a)</sup>	Spontaneous abortions	
		Number	Per cent
Less than 25	201	34	16.9
25–29	1,965	305	15.5
30–34	3,649	718	19.7
35–39	2,163	550	25.4
40–44	457	203	44.4
45 and over	20	12	60.0
Not stated	2	-	-
<b>All ages</b>	<b>8,457</b>	<b>1,822</b>	<b>21.5</b>

(a) Spontaneous abortions and pregnancies of at least 20 weeks' gestation

Table 73: Ectopic pregnancies after GIFT, 1985–1997

Outcome of pregnancy	1985–94	1995	1996	1997	1985–97
Ectopic pregnancies	242	14	11	16	283
Clinical pregnancies	6,980	654	654	505	8,793
% ectopic pregnancies	3.5	2.1	1.7	3.2	3.2
Total abortions and births	6,738	640	643	489	8,510
Ectopic pregnancy ratio <sup>(a)</sup>	1:27.8	1:45.7	1:58.5	1:30.6	1:30.1

(a) Ratio of ectopic pregnancies: total abortions and births

Table 74: Heterotopic pregnancies after GIFT, 1985–1997

Type of heterotopic pregnancy	Year							1985–97
	1985–91	1992	1993	1994	1995	1996	1997	
	<b>Number</b>							
Abortion	19	2	2	2	1	2	4	32
Birth	14	1	1	0	0	1	0	17
<b>All heterotopic pregnancies</b>	<b>33</b>	<b>3</b>	<b>3</b>	<b>2</b>	<b>1</b>	<b>3</b>	<b>4</b>	<b>49</b>
All pregnancies	4,052	1,036	1,022	870	654	654	505	8,793
	<b>Per cent</b>							
Abortion	0.5	0.2	0.2	0.2	0.2	0.3	0.8	0.4
Birth	0.3	0.1	0.1	0.0	0.0	0.2	0.0	0.2
<b>All heterotopic pregnancies</b>	<b>0.8</b>	<b>0.3</b>	<b>0.3</b>	<b>0.2</b>	<b>0.2</b>	<b>0.5</b>	<b>0.8</b>	<b>0.6</b>

Table 75: Reported complications of pregnancy, GIFT pregnancies, 1990–1997

Pregnancy complications	Number			Per cent		
	1990–95	1996	1997	1990–95	1996	1997
None	2,641	480	359	62.6	73.6	74.9
Threatened abortion	277	23	19	6.6	3.5	4.0
Antepartum haemorrhage	107	11	11	2.5	1.7	2.3
Pregnancy-induced hypertension	304	33	16	7.2	5.1	3.3
Placenta praevia	36	7	7	0.9	1.1	1.5
Other complications	852	98	67	20.2	15.0	14.0
Not stated	1,148	2	26			
<b>All pregnancies</b>	<b>5,365</b>	<b>654</b>	<b>505</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Table 76: Duration of singleton and multiple GIFT pregnancies of at least 20 weeks' gestation, 1996-1997

Gestational age (weeks)	Singleton		Twin		Triplet		All pregnancies <sup>(a)</sup>	
	Number	Per cent	Number	Per cent	Number	Per cent	Number	Per cent
20-23	6	0.9	3	1.7	2	8.3	11	1.3
24-27	5	0.7	4	2.3	3	12.5	12	1.4
28-31	7	1.0	11	6.2	4	16.7	23	2.6
32-36	66	9.8	76	42.9	15	62.5	157	17.9
37-41	582	86.5	83	46.9	-	-	665	76.0
42 or more	7	1.0	-	-	-	-	7	0.8
20 - 36	84	12.5	94	53.1	24	100.0	203	23.2
Not stated	7		-		-		7	
<b>All gestational ages</b>	<b>680</b>	<b>100.0</b>	<b>177</b>	<b>100.0</b>	<b>24</b>	<b>100.0</b>	<b>882</b>	<b>100.0</b>

(a) Includes 1 quadruplet pregnancy

Table 77: Maternal age and duration of singleton GIFT pregnancies of at least 20 weeks' gestation, 1996-1997

Gestational age (weeks)	Maternal age (years)					All ages
	Less than 25	25-29	30-34	35-39	40 and over	
	<b>Number</b>					
20-27	-	3	6	2	-	11
28-31	-	-	4	2	1	7
32-36	1	8	28	23	6	66
37 or more	5	122	236	186	40	589
Not stated	-	1	4	2	-	7
<b>All gestational ages</b>	<b>6</b>	<b>134</b>	<b>278</b>	<b>215</b>	<b>47</b>	<b>680</b>
20-36	1	11	38	27	7	84
	<b>Per cent</b>					
20-27	-	2.3	2.2	0.9	-	1.6
28-31	-	-	1.5	0.9	2.1	1.0
32-36	16.7	6.0	10.2	10.8	12.8	9.8
37 or more	83.3	91.7	86.1	87.3	85.1	87.5
<b>All gestational ages</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
20-36	16.7	8.3	13.9	12.7	14.9	12.5

Table 78: Causes of infertility and duration of singleton GIFT pregnancies of at least 20 weeks' gestation, 1996–1997

Gestational age (weeks)	Causes of infertility					All causes <sup>(a)</sup>
	Tubal	Male	Endometriosis	Multiple	Unexplained	
	<b>Number</b>					
20–27	-	2	1	3	2	11
28–31	-	-	2	1	2	7
32–36	2	3	9	14	23	66
37 or more	15	112	92	124	183	589
Not stated	1	1	1	3	1	7
<b>All gestational ages</b>	<b>18</b>	<b>118</b>	<b>105</b>	<b>145</b>	<b>211</b>	<b>680</b>
20–36	2	5	12	18	27	84
	<b>Per cent</b>					
20–27	-	1.7	1.0	2.1	1.0	1.6
28–31	-	-	1.9	0.7	1.0	1.0
32–36	11.8	2.6	8.7	9.9	11.0	9.8
37–or more	88.2	95.7	88.5	87.3	87.1	87.5
<b>All gestational ages</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
20–36	11.8	4.3	11.5	12.7	12.9	12.5

(a) Includes 83 pregnancies with 'other' or 'not stated' causes of infertility

Table 79: Plurality of GIFT pregnancies of at least 20 weeks' gestation, 1985–1997

Plurality	Number			Per cent		
	1985–95	1996	1997	1985–95	1996	1997
Singletons	4,281	391	289	74.4	75.9	78.7
Twins	1,250	108	69	21.7	21.0	18.8
Triples	205	15	9	3.6	2.9	2.5
Quadruplets	15	1	-	0.3	0.2	-
Quintuplets	2	-	-	0.0	-	-
<b>All pregnancies</b>	<b>5,753</b>	<b>515</b>	<b>367</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

**Table 80: Plurality of GIFT pregnancies of at least 20 weeks' gestation and number of oocytes transferred, 1996–1997**

Number of oocytes transferred	Plurality							
	Singleton		Twin		Triplet		All pregnancies <sup>(a)</sup>	
	Number	Per cent	Number	Per cent	Number	Per cent	Number	Per cent
1	10	83.3	2	16.7	-	-	12	100.0
2	318	82.2	65	16.8	4	1.0	387	100.0
3	306	71.2	104	24.2	19	4.4	430	100.0
4	37	86.0	6	14.0	-	-	43	100.0
5 or more	4	80.0	-	-	1	-	5	100.0
Not stated	5	-	-	-	-	-	5	-
<b>All pregnancies</b>	<b>680</b>	<b>77.1</b>	<b>177</b>	<b>20.1</b>	<b>24</b>	<b>2.7</b>	<b>882</b>	<b>100.0</b>

(a) Includes 1 quadruplet pregnancy

**Table 81: Method of delivery for singleton and multiple GIFT pregnancies of at least 20 weeks' gestation, 1996–1997**

Plurality	Method of delivery				
	Vaginal		Caesarean section		All methods <sup>(a)</sup>
	Number	Per cent	Number	Per cent	Number
Singleton	451	66.7	225	33.3	680
Twin	60	33.9	117	66.1	177
Triplet	5	20.8	19	79.2	24
Quadruplet	-	-	1	100.0	1
<b>All pregnancies</b>	<b>516</b>	<b>58.8</b>	<b>362</b>	<b>41.2</b>	<b>882</b>

(a) Includes 32 pregnancies in which the method of delivery was not stated

**Table 82: Sex of infants in singleton and multiple GIFT births of at least 20 weeks' gestation, 1985–1997**

Plurality	Male			Female			Sex ratio (M:F)		
	1985–95	1996	1997	1985–95	1996	1997	1985–95	1996	1997
	Singletons	2,216	209	152	2,045	182	135	108.4	114.8
Twins	1,266	126	72	1,225	90	66	103.3	140.0	109.1
Triplets	340	20	12	273	25	15	124.5	80.0	80.0
Quadruplets	33	2	-	27	2	-	122.2	100.0	-
Quintuplets	4	-	-	6	-	-	66.7	-	-
<b>All births</b>	<b>3,859</b>	<b>357</b>	<b>236</b>	<b>3,576</b>	<b>299</b>	<b>216</b>	<b>107.9</b>	<b>119.4</b>	<b>109.3</b>

Note. Infant's sex was not stated or indeterminate for 33 births

**Table 83: Birthweight of GIFT live births and stillbirths, 1996–1997**

Birthweight (g)	Live births		Stillbirths		All births <sup>(a)</sup>	
	Number	Per cent	Number	Per cent	Number	Per cent
Less than 500	6	0.6	8	72.7	14	1.3
500–999	17	1.6	2	18.2	22	2.0
1000–1499	48	4.6	-	0.0	50	4.5
1500–1999	80	7.6	-	0.0	83	7.5
2000–2499	152	14.5	1	9.1	156	14.2
2500–2999	252	24.1	-	0.0	258	23.4
3000–3499	280	26.7	-	0.0	294	26.7
3500–3999	162	15.5	-	0.0	171	15.5
4000 and over	50	4.8	-	0.0	54	4.9
Not stated	2		6		8	
<b>All birthweights</b>	<b>1,049</b>	<b>100.0</b>	<b>17</b>	<b>100.0</b>	<b>1,110</b>	<b>100.0</b>
Mean birthweight (g)	2,842		594		2,824	

(a) Includes 44 infants with unknown outcome

**Table 84: Birthweight of infants in singleton and multiple GIFT births of at least 20 weeks' gestation, 1996–1997**

Birthweight (g)	Singleton		Twin		Triplet		All births <sup>(a)</sup>	
	Number	Per cent	Number	Per cent	Number	Per cent	Number	Per cent
Less than 500	4	0.6	6	1.7	4	5.6	14	1.3
500–999	4	0.6	7	2.0	10	14.1	22	2.0
1000–1499	9	1.3	24	6.8	14	19.7	50	4.5
1500–1999	10	1.5	43	12.2	30	42.3	83	7.5
2000–2499	38	5.6	106	30.0	12	16.9	156	14.2
2500–2999	140	20.8	118	33.4	-	-	258	23.4
3000–3499	252	37.4	41	11.6	1	1.4	294	26.7
3500–3999	163	24.2	8	2.3	-	-	171	15.5
4000 and over	54	8.0	-	-	-	-	54	4.9
Less than 2500	65	9.6	186	52.7	70	98.6	325	29.5
Not stated	6		1		1		8	
<b>All birthweights</b>	<b>680</b>	<b>100.0</b>	<b>354</b>	<b>100.0</b>	<b>72</b>	<b>100.0</b>	<b>1,110</b>	<b>100.0</b>
Mean birthweight (g)	3,209		2,367		1,532		2,824	

(a) Includes 4 quadruplet births



**Table 85: Outcome of infants in singleton and multiple GIFT births of at least 20 weeks' gestation, 1996-1997**

Outcome	Singleton	Twin	Triplet	Quadruplet	All births <sup>(a)</sup>
Live births	673	348	68	4	1,093
Stillbirths	7	6	4	-	17
<b>All births</b>	<b>680</b>	<b>354</b>	<b>72</b>	<b>4</b>	<b>1,110</b>
Neonatal deaths	1	4	5	-	10
Perinatal deaths	8	10	9	-	27
Stillbirth rate per 1,000 total births	10.3	16.9	55.6	-	15.3
Neonatal death rate per 1,000 live births	1.5	11.5	73.5	-	9.1
Perinatal mortality rate per 1,000 total births	11.8	28.2	125.0	-	24.3

(a) Live births include 44 births for which birth status was not recorded

**Table 86: Major congenital malformations in singleton and multiple GIFT births of at least 20 weeks' gestation, 1985-1997**

Outcome	Singleton	Multiple	All births <sup>(a)</sup>
Total births	4,991	3,617	8,608
Congenital malformations			
— number	137	72	209
— rate (per cent)	2.7	2.0	2.4

(a) Includes 32 abortions for fetal abnormality at gestational ages of at least 16 weeks

## 6 Bibliography

This bibliography lists references to scientific, epidemiological and social studies on in-vitro fertilisation and other methods of assisted conception in Australia and New Zealand which have been published or made available since the previous report for 1996.

1. Almahbobi G, Misajon A, Hutchinson P, Lolatgis N, & Trounson AO 1998. Hyperexpression of epidermal growth factor receptors in granulosa cells from women with polycystic ovary syndrome. *Fertility & Sterility* 70(4):750–8.
2. Baker HW 1997. Male infertility of undetermined etiology. [Review] [4 refs] *Current Therapy in Endocrinology & Metabolism* 6:359–63.
3. Baker HW 1998. Marvellous ICSI: the viewpoint of a clinician. *International Journal of Andrology* 21(5):249–52.
4. Baker HW 1997. Medical treatment for idiopathic male infertility: is it curative or palliative? [Review] [63 refs] *Baillieres Clinical Obstetrics & Gynaecology* 11(4):673–89.
5. Bourne H, Stern K, Clarke G, Pertile M, Speirs A & Baker HW 1997. Delivery of normal twins following the intracytoplasmic injection of spermatozoa from a patient with 47,XXY Klinefelter's syndrome. *Human Reproduction* 12(11):2447–50.
6. Bowman M, De Boer K, Cullinan R, Catt J & Jansen R 1998. Do alterations in the sex ratio occur at fertilization? A case report using fluorescent in situ hybridization. *Journal of Assisted Reproduction & Genetics* 15(5):320–2.
7. Clarke GN 1997. A simple monthly means chart system for monitoring sperm concentration. *Human Reproduction* 12(12):2710–2.
8. Clarke GN, Garrett C & Baker HW 1998. Quantitative sperm mucus penetration: modified formulae for calculating penetration efficiency. *Human Reproduction* 13(5):1255–9.
9. Cummins JM, Jequier AM, Martin R, Mehmet D & Goldblatt J 1998. Semen levels of mitochondrial DNA deletions in men attending an infertility clinic do not correlate with phenotype. *International Journal of Andrology* 21(1):47–52.
10. Cummins JM, Wakayama T & Yanagimachi R 1997. Fate of microinjected sperm components in the mouse oocyte and embryo. *Zygote* 5(4):301–8.
11. Daniels K 1997. The controversy regarding privacy versus disclosure among patients using donor gametes in assisted reproductive technology. *Journal of Assisted Reproduction & Genetics* 14(7):373–5.

12. Daniels KR & Burn I 1997. Access to assisted human reproduction services by minority groups. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 37(1):79–85.
13. Daniels KR & Hall DJ 1997. Semen donor recruitment strategies – a non-payment based approach. [Review] [39 refs] *Human Reproduction* 12(10):2330–5.
14. Daniels KR, Lewis GM & Curson R 1997. Information sharing in semen donation: the views of donors. *Social Science & Medicine* 44(5):673–80.
15. De Jonge CJ, Flaherty SP, Barnes AM, Swann NJ & Matthews CD 1997. Failure of multitube sperm swim-up for sex preselection. *Fertility & Sterility* 67(6):1109–14.
16. Downie SE, Flaherty SP & Matthews CD 1997. Detection of chromosomes and estimation of aneuploidy in human spermatozoa using fluorescence in-situ hybridization. [Review] [108 refs] *Molecular Human Reproduction* 3(7):585–98.
17. Downie SE, Flaherty SP, Swann NJ & Matthews CD 1997. Estimation of aneuploidy for chromosomes 3, 7, 16, X and Y in spermatozoa from 10 normospermic men using fluorescence in-situ hybridization. *Molecular Human Reproduction* 3(9):815–9.
18. Driscoll GL, Tyler JP & Carpenter D 1997. Variation in the determination of follicular diameter: an inter-unit pilot study using an ultrasonic phantom. *Human Reproduction* 12(11):2465–8.
19. Driscoll GL, Tyler JP, Knight DC, Cooke S, Kime L, Clark L & Bernstein J 1998. Failure to collect oocytes in assisted reproductive technology: a retrospective. *Human Reproduction* 13(1):84–7.
20. Edirisinghe WR, Murch A, Junk S & Yovich JL 1997. Cytogenetic abnormalities of unfertilized oocytes generated from in-vitro fertilization and intracytoplasmic sperm injection: a double-blind study. *Human Reproduction* 12(12):2784–91.
21. Edirisinghe WR, Murch AR, Junk SM & Yovich JL 1998. Cytogenetic analysis of unfertilized oocytes following intracytoplasmic sperm injection using spermatozoa from a globozoospermic man. *Human Reproduction* 13(11):3094–8.
22. Eldar-Geva T, Lowe PJ, MacLachlan V, Rombauts L & Healy DL 1998. Different influence of incongruent follicular development on in vitro fertilization-embryo transfer and gamete intrafallopian transfer pregnancy rates. *Fertility & Sterility* 70(6):1039–43.
23. Eldar-Geva T, Meagher S, Healy DL, MacLachlan V, Breheny S & Wood C 1998. Effect of intramural, subserosal, and submucosal uterine fibroids on the outcome of assisted reproductive technology treatment. *Fertility & Sterility* 70(4):687–91.
24. Flaherty SP, Michalowska J, Swann NJ, Dmowski WP, Matthews CD & Aitken RJ 1997. Albumin gradients do not enrich Y-bearing human spermatozoa. *Human Reproduction* 12(5):938–42.

25. Flaherty SP, Payne D & Matthews CD 1998. Fertilization failures and abnormal fertilization after intracytoplasmic sperm injection. *Human Reproduction* 13 Suppl 1:155–64.
26. Fong CY, Bongso A, Ng SC, Anandakumar C, Trounson A & Ratnam S 1997. Ongoing normal pregnancy after transfer of zona-free blastocysts: implications for embryo transfer in the human. *Human Reproduction* 12(3):557–60.
27. Gardner DK & Lane M 1997. Culture and selection of viable blastocysts: a feasible proposition for human IVF? [Review] [139 refs] *Human Reproduction Update* 3(4): 367–82.
28. Gardner DK, Vella P, Lane M, Wagley L, Schlenker T & Schoolcraft WB 1998. Culture and transfer of human blastocysts increases implantation rates and reduces the need for multiple embryo transfers. *Fertility & Sterility* 69(1):84–8.
29. Gibson FL, Ungerer JA, Leslie GI, Saunders DM & Tennant CC 1998. Development, behaviour and temperament: a prospective study of infants conceived through in-vitro fertilization. *Human Reproduction* 13(6):1727–32.
30. Gook D, Osborn SM, Bourne H, Edgar DH & Speirs AL 1998. Fluorescent study of chromatin and tubulin in apparently unfertilized human oocytes following ICSI. *Molecular Human Reproduction* 4(12):113–5.
31. Gomez MC, Catt JW, Gillan L, Evans G & Maxwell WM 1997. Effect of culture, incubation and acrosome reaction of fresh and frozen-thawed ram spermatozoa for in vitro fertilization and intracytoplasmic sperm injection. *Reproduction, Fertility, & Development* 9(7):665–73.
32. Healy DL 1997. Occult ovarian failure. [Review] [6 refs] *Current Therapy in Endocrinology & Metabolism* 6:226–7.
33. Holden CA, Fuscaldo GF, Jackson P, Cato A, Southwick GJ, Hauser R, Temple-Smith PD & McLachlan RI 1997. Frozen-thawed epididymal spermatozoa for intracytoplasmic sperm injection. *Fertility & Sterility* 67(1):81–7.
34. Horne G, Critchlow JD, Newman MC, Edozien L, Matson PL & Lieberman BA 1997. A prospective evaluation of cryopreservation strategies in a two-embryo transfer programme. *Human Reproduction* 12(3):542–7.
35. Imade GE, Baker HW, de Kretser DM & Hedger MP 1997. Immunosuppressive activities in the seminal plasma of infertile men: relationship to sperm antibodies and autoimmunity. *Human Reproduction* 12(2):256–62.
36. Jansen RP 1997. Evidence-based ethics and the regulation of reproduction. [Review] [39 refs] *Human Reproduction* 12(9):2068–75.
37. Jansen RP 1997. Reproductive medicine and the social state of childlessness. *Medical Journal of Australia* 167(6):321–3.

38. Jequier AM 1998. Clinical andrology today [letter]. *Human Reproduction* 13(7):2036–7.
39. Jequier AM 1997. Clinical assessment of male infertility in the era of intracytoplasmic sperm injection. [Review] [59 refs] *Baillieres Clinical Obstetrics & Gynaecology* 11(4):617–39.
40. Jequier AM 1998. Vasectomy related infertility: a major and costly medical problem. *Human Reproduction* 13(7):1757–9.
41. Joesbury KA, Edirisinghe WR, Phillips MR & Yovich JL 1998. Evidence that male smoking affects the likelihood of a pregnancy following IVF treatment: application of the modified cumulative embryo score. *Human Reproduction* 13(6):1506–13.
42. Johnson N, McComb P & Gudex G 1998. Heterotopic pregnancy complicating in vitro fertilization. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 38(2): 151–5.
43. Jones GM, Trounson AO, Gardner DK, Kausche A, Lolatgis N & Wood C 1998. Evolution of a culture protocol for successful blastocyst development and pregnancy. *Human Reproduction* 13(1):169–77.
44. Jones GM, Trounson AO, Lolatgis N & Wood C 1998. Factors affecting the success of human blastocyst development and pregnancy following in vitro fertilization and embryo transfer. *Fertility & Sterility* 70(6):1022–9.
45. Lane M & Gardner DK 1997. Nonessential amino acids and glutamine decrease the time of the first three cleavage divisions and increase compaction of mouse zygotes in vitro. *Journal of Assisted Reproduction & Genetics* 14(7):398–403.
46. Leslie GI, Gibson FL, McMahon C, Tennant C & Saunders DM 1998. Infants conceived using in-vitro fertilization do not over-utilize health care resources after the neonatal period. *Human Reproduction* 13(8):2055–9.
47. Liu DY & Baker HW 1998. Calcium ionophore-induced acrosome reaction correlates with fertilization rates in vitro in patients with teratozoospermic semen. *Human Reproduction* 13(4):905–10.
48. Liu DY, Baker HW, Pearse MJ & d'Apice AJ 1997. Normal sperm-zona pellucida interaction and fertilization in vitro in alpha-1, 3-galactosyltransferase gene knockout mice [letter]. *Molecular Human Reproduction* 3(11):1015–6.
49. Liu DY & Baker HW 1997. Protein kinase C plays an important role in the human zona pellucida-induced acrosome reaction. *Molecular Human Reproduction* 3(12):1037–43.
50. Lopata A, Oliva K, Stanton PG & Robertson DM 1997. Analysis of chorionic gonadotrophin secreted by cultured human blastocysts. *Molecular Human Reproduction* 3(6):517–21.

51. Lowe PJ, Mamers PM, Sturrock TV & Healy D 1998. A casemix cost comparison of 2 treatments for ectopic pregnancy. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 38(3):333-5.
52. Martini E, Flaherty SP, Swann NJ, Payne D & Matthews CD 1997. Analysis of unfertilized oocytes subjected to intracytoplasmic sperm injection using two rounds of fluorescence in-situ hybridization and probes to five chromosomes. *Human Reproduction* 12(9):2011-8.
53. Matson PL 1997. Clinical value of tests for assessing male infertility. [Review] [90 refs] *Baillieres Clinical Obstetrics & Gynaecology* 11(4):641-54.
54. Mortimer ST 1997. A critical review of the physiological importance and analysis of sperm movement in mammals. [Review] [346 refs] *Human Reproduction Update* 3(5):403-39.
55. Mortimer ST, Swan MA & Mortimer D 1998. Effect of seminal plasma on capacitation and hyperactivation in human spermatozoa. *Human Reproduction* 13(8):2139-46.
56. Munne S, Magli C, Adler A, Wright G, de Boer K, Mortimer D, Tucker M, Cohen J & Gianaroli L 1997. Treatment-related chromosome abnormalities in human embryos. *Human Reproduction* 12(4):780-4.
57. Naysmith TE, Blake DA, Harvey VJ & Johnson NP 1998. Do men undergoing sterilizing cancer treatments have a fertile future? *Human Reproduction* 13(11):3250-5.
58. Oehninger S, Pike I & Slotnick N 1998. Hemochromatosis and male infertility. *Obstetrics & Gynecology* 92(4 Pt 2):652-3.
59. O'Neill C 1997. Evidence for the requirement of autocrine growth factors for development of mouse preimplantation embryos in vitro. *Biology of Reproduction* 56(1):229-37.
60. Payne D, Flaherty SP, Barry MF & Matthews CD 1997. Preliminary observations on polar body extrusion and pronuclear formation in human oocytes using time-lapse video cinematography. *Human Reproduction* 12(3):532-41.
61. Rombauts L, Suikkari AM, MacLachlan V, Trounson AO & Healy DL 1998. Recruitment of follicles by recombinant human follicle-stimulating hormone commencing in the luteal phase of the ovarian cycle. *Fertility & Sterility* 69(4):665-9.
62. Sathananthan AH, Szell A, Ng SC, Kausche A, Lacham-Kaplan O & Trounson A 1997. Is the acrosome reaction a prerequisite for sperm incorporation after intra-cytoplasmic sperm injection (ICSI)? *Reproduction, Fertility & Development* 9(7):703-9.
63. Sathananthan AH, Tatham B, Dharmawardena V, Grills B, Lewis I & Trounson A 1997. Inheritance of sperm centrioles and centrosomes in bovine embryos. *Archives of Andrology* 38(1):37-48.

64. Shaw J & Trounson A 1997. Oncological implications in the replacement of ovarian tissue. [Review] [16 refs] *Human Reproduction* 12(3):403-5.
65. Speirs AL 1998. Aspects of epigenetic factors in artificial procreation [comment]. *Human Reproduction* 13(4):787-8.
66. Stern C, Chamley L, Hale L, Kloss M, Speirs A & Baker HW 1998. Antibodies to beta2 glycoprotein I are associated with in vitro fertilization implantation failure as well as recurrent miscarriage: results of a prevalence study. *Fertility & Sterility* 70(5):938-44.
67. Swan MA & Alboghobeish N 1997. Improved preservation of the ram spermatozoan plasma membrane using betaine in the primary fixative. *Journal of Microscopy* 187(3):167-9.
68. Swan MA 1997. Improved preservation of ultrastructural morphology in human spermatozoa using betaine in the primary fixative. *International Journal of Andrology* 20(1):45-54.
69. Tadokoro N, Vollenhoven B, Clark S, Baker G, Kovacs G, Burger H & Healy D 1997. Cumulative pregnancy rates in couples with anovulatory infertility compared with unexplained infertility in an ovulation induction programme. *Human Reproduction* 12(9):1939-44.
70. Trounson A, Anderiesz C, Jones GM, Kausche A, Lolatgis N & Wood C 1998. Oocyte maturation. [Review] [13 refs] *Human Reproduction* 13 Suppl 3:52-62; discussion 71-5.
71. Trounson A, Pera M 1998. Potential benefits of cell cloning for human medicine. [Review] [30 refs] *Reproduction, Fertility & Development* 10(1):121-5.
72. Trounson AO 1997. Cloning: potential benefits for human medicine [editorial]. [Review] [14 refs] *Medical Journal of Australia* 167(11-12):568-9.
73. Wood CE, Shaw JM & Trounson AO 1997. Cryopreservation of ovarian tissue: Potential "reproductive insurance" for women at risk of early ovarian failure. [Review] [50 refs] *Medical Journal of Australia* 166(7):366-9.
74. Woolcott R & Stanger J 1998. Ultrasound tracking of the movement of embryo-associated air bubbles on standing after transfer. *Human Reproduction* 13(8):2107-9.

# Appendix 1: Definitions

**Clinical pregnancy:** Any type of pregnancy except that diagnosed only by measuring levels of human chorionic gonadotrophin. This definition includes ectopic pregnancy, blighted ovum and spontaneous abortion.

**Conception cohort:** A designated group of pregnancies resulting from conception in a specified period of time (usually either a single year or several years combined).

**Ectopic pregnancy:** Pregnancy occurring outside the uterus.

**Fetal death (stillbirth):** Death prior to the complete expulsion or extraction from its mother of a product of conception of 20 or more completed weeks of gestation or of 400 g or more birthweight; the death is indicated by the fact that after such separation the fetus does not breathe or show any other evidence of life, such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles.

**Heterotopic pregnancy:** Heterotopic pregnancies are those in which there is both a uterine and tubal (ectopic) pregnancy simultaneously. The uterine pregnancy may abort or may continue on to a birth.

**Live birth:** Infant with signs of life after pregnancy of at least 20 weeks' gestation.

**Live-birth pregnancy:** A pregnancy resulting in one or more live births.

**Low birthweight:** A liveborn or stillborn infant weighing less than 2500 g at birth.

**Neonatal death:** A death of a liveborn infant within 28 days of birth (expressed as a rate per 1,000 live births).

**Perinatal death:** Includes both stillbirths and neonatal deaths (expressed as a rate per 1,000 total births).

**Postneonatal death:** A death of a liveborn infant more than 28 days after birth but within the first year (expressed as a rate per 1,000 live births).

**Preterm birth:** A liveborn or stillborn infant of less than 37 weeks' gestation.

**Spontaneous abortion:** Pregnancy detected clinically or by ultrasound and less than 20 weeks' gestation (from the first day of the last menstrual period).

**Stillbirth:** See fetal death.

**Viable pregnancy:** A pregnancy of at least 20 weeks' gestation.



# Appendix 2: Notification form

AIHW NATIONAL PERINATAL STATISTICS UNIT / FERTILITY SOCIETY OF AUSTRALIA

## REGISTER OF PREGNANCIES AFTER IVF OR RELATED PROCEDURES

Please complete all data items by ticking relevant boxes

IVF Unit/Hospital: \_\_\_\_\_ Identification number: \_\_\_\_\_

Usual home address Suburb/Town _____	: Marital status : [ ] Married/De facto	: Date of birth : Mother ___/___/___	Age ___ yrs
State _____ Postcode _____	: [ ] Single [ ] Other	: Father ___/___/___	___ yrs
NUMBER OF PREVIOUS PREGNANCIES: Current marriage _____ Previous marriages _____		TYPE OF CONCEPTION IN CURRENT PREGNANCY:	
Livebirths _____	Mother:livebirths _____	[ ] IVF	[ ] PROST/ZIFT [ ] TEST
Abortions _____	other _____	[ ] GIFT	[ ] ICSI [ ] SUZI
Other _____	Father:livebirths _____	[ ] Epididymal sperm	[ ] Assisted hatching
	other _____	[ ] Other (specify) _____	
Did this pregnancy result from use of:			
[ ] Donor sperm	[ ] Donor oocyte	[ ] Frozen embryo	
[ ] Donor embryos	[ ] Frozen oocytes		
What was the date of embryo freezing? ---/---/--- : If donor oocyte or embryo, what was : the age of the donor?			
What was the date of embryo transfer? ___/___/___ : _____ yrs			
CAUSE OF INFERTILITY PRIOR TO THIS PREGNANCY [ ] Unknown cause			
Tubal [ ] Tubal obstruction	[ ] Previous ectopic	[ ] Salpingectomy	
[ ] Sterilization	[ ] Pelvic adhesions	[ ] Pelvic inflammatory disease	
[ ] Other tubal (specify) _____			
Male factor [ ] Azoospermia	[ ] Oligospermia		
[ ] Increased abnormal sperm	[ ] Male sperm antibodies		
[ ] Decreased motility	[ ] Other male (specify) _____		
[ ] Endometriosis	[ ] Ovulation defects	[ ] Maternal sperm antibodies	
[ ] 'Hostile' cervical mucus	[ ] Other cause (specify) _____		
DURATION OF INFERTILITY (before first IVF/GIFT pregnancy) _____ years			
DRUGS USED TO INDUCE OVULATION IN OOCYTE RETRIEVAL CYCLE (specify each separately)			
[ ] Clomiphene	[ ] hMG	[ ] hCG	[ ] Endogenous LH surge
[ ] FSH	[ ] Recomb DNA FSH	[ ] Recomb DNA LH	
[ ] GnRH-agonist (specify) _____	[ ] short protocol	[ ] long protocol	(previous luteal phase)
[ ] None	[ ] Other (specify) _____		
DRUGS USED DURING CYCLE IN WHICH FROZEN EMBRYOS WERE TRANSFERRED(specify each separately)			
[ ] None	[ ] Oestrogen/progesterone	[ ] Other (specify) _____	

SPECIFY IN WHICH OOCYTE RETRIEVAL CYCLE THE PREGNANCY OCCURRED _____		METHOD OF COLLECTING OOCYTES	
Number of oocytes collected _____		[ ] Laparoscopy	
IF DONOR OOCYTES WERE USED, IN WHICH INDUCTION CYCLE DID PREGNANCY OCCUR _____		[ ] Ultrasound-guided transvaginal	
		[ ] Other (specify) _____	
Date of fertilization (or GIFT, etc) --/--/--		Number of embryos/ova transferred _____	
Was the patient hospitalised for ovarian hyperstimulation syndrome? [ ] Yes [ ] No			
DRUGS USED IN LUTEAL PHASE		OBSTETRIC COMPLICATIONS	
[ ] hCG: specify dose and duration _____		[ ] None [ ] Pregnancy-induced hypertension	
[ ] Progesterone: specify dose and duration _____		[ ] Threatened abortion	
[ ] Oestrogen/progesterone (frozen embryo transfer) _____		[ ] Placenta praevia	
[ ] Other (specify) _____		[ ] Antepartum haemorrhage	
[ ] None		[ ] Embryo reduction	
		[ ] Other _____	
NUMBER OF SACS SEEN IN EARLY PREGNANCY ON ULTRASOUND EXAMINATION _____		[ ] Ultrasound not done	
PREGNANCY OUTCOME		[ ] Ectopic pregnancy	
[ ] Spontaneous abortion (date __/__/__)		[ ] Ovarian pregnancy	
[ ] Missed abortion (date of curette --/--/--)		[ ] Blighted ovum	
[ ] Induced abortion (date __/__/__, specify malformations _____)			
[ ] Other (e.g. combined pregnancy) _____			
[ ] Pregnancy of 20 weeks or more		Date of birth __/__/__	
[ ] Multiple births (number _____)			
METHOD OF DELIVERY		[ ] Vaginal [ ] Caesarean section	
LIVEBIRTHS AND STILLBIRTHS	1	2	3
Sex	M F	M F	M F
Birthweight	_____g	_____g	_____g
Condition at birth (delete one)	Live birth/ Stillbirth	Live birth/ Stillbirth	Live birth/ Stillbirth
If baby died, date of death	__/__/__	__/__/__	__/__/__
Any congenital malformations?	[ ] Yes [ ] No	[ ] Yes [ ] No	[ ] Yes [ ] No
Specify malformations or other abnormalities			