
2022 SAMPLE CHARACTERISTICS



In 2022, 100 participants, recruited from Perth, WA were interviewed.



The median age in 2022 was 21, and 71% identified as male.




In the 2022 sample, 37% were enrolled students, 38% were employed full time and 15% were unemployed.



Participants were recruited on the basis that they had consumed ecstasy and/or other illicit stimulants at least monthly in the past 6 months.

PAST 6 MONTH USE OF SELECTED DRUGS



MENTAL HEALTH AND SEXUAL HEALTH BEHAVIOURS




MH issue

32%

Seen a MH professional

In the total sample, 57% self-reported a mental health issue and 32% had seen a mental health professional in the past 6 months.



Of those who had a mental health condition, the three most common mental health issues reported were anxiety (62%), depression (50%) and any personality disorder (12%).




sexual activity

34%


Had an STI test

In the total sample, 76% reported sexual activity in the past 4 weeks, and 34% had a sexual health check in the past 6 months.



Sexual risk behaviours among those who reported any sexual activity in the past four weeks and were able to comment

OTHER RISK BEHAVIOURS




Among recent drivers, 55% reported driving a vehicle within 3 hours of consuming illicit drugs and 35% while over the legal limit of alcohol.



31% 2022

In the 2022 sample, 31% reported a non-fatal depressant overdose in the previous 12 months, an increase relative to 2021 (17%).



In the total sample, 87% reported concurrent use of two or more substances on the last occasion of ecstasy or related drug use.




depressants

9% Stimulants and Stimulants, depressants and


cannahis

The most commonly used combinations of drug classes on the last occasion of ecstasy or related drua use.

ECSTASY



Past 6 month use of ecstasy capsules, crystal, pills, and powder in 2022.



Of those who had recently consumed ecstasy, 10% reported weekly or more frequent use.



2 Capsules 2 Pills



0.30 grams of crystal 0.30 grams of powder

Median amounts of ecstasy consumed in a 'typical' session using each form.




In 2022, more participants perceived the availability of the powder form of ecstasy as 'difficult' relative to 2021.


METHAMPHETAMINE




Past 6 month use of any methamphetamine, crystal, powder and base in 2022.



Of those who had recently consumed methamphetamine, small numbers reported weekly or more frequent use (n≤5).




91% of participants who had recently used crystal smoked it.




Of those who could comment, 86% perceived crystal methamphetamine to be 'easy' or 'very easy' to obtain.

COCAINE




Past 6 month use of any cocaine remained stable between 2021 and 2022.



Of those who had recently consumed cocaine, small numbers reported weekly or more frequent use (n≤5).




In 2022, the median price of a gram of cocaine remained stable at \$400.




Of those who could comment, 62% perceived cocaine to be 'easy' or 'very easy' to obtain.


CANNABIS AND/OR CANNABINOID REL




Past 6 month use of non-prescribed cannabis and/ or cannabinoid related products remained stable between 2021 and 2022.



Of those who had recently consumed non-prescribed cannabis and/or cannabinoid related products, 63% reported weekly or more frequent use.



Of participants who had consumed non-prescribed cannabis and/or cannabinoid related products in the last 6 months, 94% had smoked it.



Of those who could comment, 95% perceived hydro to be 'easy' or 'very easy' to obtain.